LMU–ASC 56/06
MPP–2006–110
DISTA–UPO–06
hep-th/0609013

arXiv:hep-th/0609013v2 16 Nov 2006

Moduli Stabilization in Type *IIB* Orientifolds (II)

D. Lüst[a,b], S. Reffert[b], E. Scheidegger[c], W. Schulgin[b] and S. Stieberger[a]

[a] *Arnold–Sommerfeld–Center for Theoretical Physics,*
Department für Physik, Ludwig–Maximilians–Universität München,
Theresienstraße 37, 80333 München, Germany

[b] *Max–Planck–Institut für Physik,*
Föhringer Ring 6, 80805 München, Germany

[c] *Universita del Piemonte Orientale "A. Avogadro",*
Dip. Scienze e Tecnologie Avanzate,
Via Bellini, 25/g, 15100 Alessandria, Italy
INFN – Sezione di Torino, Italy

Abstract

We discuss general properties of moduli stablization in KKLT scenarios in type IIB orientifold compactifications X_6. In particular, we find conditions for the Kähler potential to allow a KKLT scenario for a manifold X_6 without complex structure moduli, *i.e.* $h^{(-)}_{(2,1)}(X_6) = 0$. This way, a whole class of type IIB orientifolds with $h^{(-)}_{(2,1)}(X_6) = 0$ is ruled out. This excludes in particular all $\mathbf{Z}_N$–and $\mathbf{Z}_N \times \mathbf{Z}_M$–orientifolds X_6 with $h_{(2,1)}(X_6) = 0$ for a KKLT scenario. This concerns $\mathbf{Z}_3, \mathbf{Z}_7, \mathbf{Z}_3 \times \mathbf{Z}_3, \mathbf{Z}_4 \times \mathbf{Z}_4, \mathbf{Z}_6 \times \mathbf{Z}_6$ and $\mathbf{Z}_2 \times \mathbf{Z}_{6'}$ –both at the orbifold point and away from it. Furthermore, we propose a mechanism to stabilize the Kaehler moduli accociated to the odd cohomology $H^{(1,1)}_-(X_6)$.

In the second part of this work we discuss the moduli stabilization of resolved type IIB $\mathbf{Z}_N$– or $\mathbf{Z}_N \times \mathbf{Z}_M$–orbifold/orientifold compactifications. As examples for the resolved $\mathbf{Z}_6$ and $\mathbf{Z}_2 \times \mathbf{Z}_4$ orbifolds we fix all moduli through a combination of fluxes and racetrack superpotential.

Contents

1. Introduction 2

2. Moduli stabilization and stability in type IIB orientifolds 6

  2.1. Calabi–Yau orientifolds of type IIB with $D3/D7$–branes 6

  2.2. Stabilization of Kähler moduli associated to the cohomology $H_+^{(1,1)}(X_6)$ . . . 9

  2.3. (In)stability of flux compactifications without complex structure moduli . . . 12

  2.4. Examples 22

3. Stabilization of Kähler moduli associated to the cohomology $H_-^{(1,1)}(X_6)$ 23

4. Instanton effects 29

  4.1. Calculation of χ_{D3} for divisors with $h_{(0,0)}=1$, $h_{(0,1)}=h_{(0,2)}=0$ 30

  4.2. General case: $h_{(0,1)}$, $h_{(0,2)}\neq 0$ 32

5. Fixing all moduli in resolved type IIB $\mathbf{Z}_N$– and $\mathbf{Z}_N\times\mathbf{Z}_M$–orientifolds 34

  5.1. Type IIB orientifolds of $\mathbf{Z}_N$– and $\mathbf{Z}_N\times\mathbf{Z}_M$–orbifolds 34

  5.2. Resolved toroidal orientifolds as candidate models for a KKLT scenario . . . 37

  5.3. Complex structure and dilaton stabilization through 3–form flux 40

  5.4. Kähler moduli stabilization 46

  5.5. Orbifolds without complex structure moduli 57

6. Concluding remarks 60

Appendix A. Complex structures of $\mathbf{Z}_N$– and $\mathbf{Z}_N\times\mathbf{Z}_M$–orbifolds 63

1. Introduction

As it is known already for some time, the landscape of consistent string compactifications is vast and contains many possible solutions. In particular, a large class of lower-dimensional string vacua is formed by four–dimensional, supersymmetric string compactifications containing massless scalar fields with arbitrary vacuum expectation values in the low energy effective field theory. These fields are in one-to-one correspondence with the continuous parameters of the underlying background geometry, and hence they are called moduli fields. However, massless moduli are unacceptable from the phenomenological point of view, since such fields give rise to a fifth force of about gravitational strength. Hence mechanisms for stabilizing moduli in string theory must be explored. Getting rid of moduli means that, seen from a stringy or geometrical point of view, we are looking for internal conformal field theories resp. for background spaces which are rigid, *i.e.* do not allow any continuous deformations. On the other hand, within the effective field theory, moduli stabilization is achieved by creating an effective potential that lifts all flat directions of the scalar fields.

To arrive at the goal of constructing string vacua with a tiny, positive de Sitter (dS) cosmological constant, $\Lambda_{dS}/M^4_{\text{Planck}} \sim 10^{-120}$, and without massless moduli, the following two-step procedure, proposed by KKLT in the context of type type IIB orientifold compactifications, seems to be a promising scenario [1]: in the first step one constructs compactifications that lead to four-dimensional, N=1 supersymmetric anti–de Sitter (AdS) string vacua with all moduli fixed in the minimum of the F-term effective scalar potential. Here the value of the negative AdS cosmological constant is of the order of the phenomenologically required scale for supersymmetry breaking in the supersymmetric Standard Model (SSM), *i.e.* $|\Lambda_{AdS}| \sim \mathcal{O}(1TeV)^4$. This method has the advantage that supersymmetric AdS-vacua with stabilized moduli are much easier to be constructed than supersymmetric Minkowski vacua with stabilized moduli fields. The second step of the KKLT scenario is to break supersymmetry by uplifting the AdS minimum to a dS vacuum with a tiny positive cosmological constant. Possible mechanisms for this uplift are either an explicit supersymmetry breaking by adding anti D-branes or spontaneous supersymmetry breaking by additional D-term contributions [2] to the effective scalar potential. Although many details of the uplift procedure still have to be worked out in concrete models, this proposal is attractive since the observed supersymmetry breaking scale in the SSM, like the gravitino mass, is set already by the first step of the KKLT procedure, and hence is robust against the details of the uplift. The second virtue of this scheme is that the masses of the moduli are also quite insensitive against many details of the uplift, provided that the $(\text{mass})^2$ matrix M^2_{ij} of the moduli is positive definite already in the AdS minimum of the effective potential, and that its eigenvalues are large compared to $|\Lambda_{AdS}|$. As we will see, the requirement of positive definiteness of M^2_{ij} can fail in explicit string compactifications.

On the other hand, the KKLT scenario requires a certain amount of fine tuning of parameters, in particular to achieve the necessary hierarchy of scales $\Lambda_{dS} \ll |\Lambda_{AdS}| \ll M_{\rm Planck}$. Taking into account the huge number of points in the string landscape, this hierarchy can possibly find a statistical or even an anthropic explanation.

In this paper we continue our previous work [3] on type IIB orientifold compactifications, here addressing the question for which concrete six–dimensional Calabi-Yau spaces all moduli can be stabilized in the large radius limit. Concretely we will investigate type IIB orientifolds on six–dimensional toroidal orbifolds away from the singular orbifold point, *i.e.* we will consider the case where the orbifold singularities are resolved by blow–ups. Our choice of background geometries is dictated by a kind of a compromise: on the one hand, in order to get a concrete handle on the KKLT scenario, we need to investigate models where calculability of the effective action after compactification is still possible due to the orbifold geometry. On the other hand, these background spaces are in fact smooth Calabi–Yau manifolds with a lot of non–trivial properties, such as a relatively large number of moduli fields. A detailed investigation of the properties of these spaces is performed in a companion paper [4]. A preview of some of these results has been given in [5].

In addition, it was independently demonstrated that type IIB orientifolds on orbifolds have several attractive phenomenological features, such as a non-vanishing probability to find the SSM in the open string spectrum on D–branes of the theory (for the $\mathbf{Z}_2 \times \mathbf{Z}_2$ orientifolds this probability is about one in a billion [6]). In contrast to the observable gauge and matter sectors of the SSM, the question of moduli stabilization on the other hand focuses on the properties of the hidden sector, which is coupled to the SSM sector through gravitational interactions or also through new gauge interactions. Hence the SSM sector and hidden sector can be regarded as two different modules which eventually have to be implanted at the same time at two different corners of the internal CY space. To analyze whether this is possible is beyond the scope of this paper. In this work we limit ourselves moreover to the stabilization of the closed string moduli of the background.

The successful implementation of the of the KKLT scenario requires that the following requirements are met in concrete CY compactifications:

(i) Turning on 3–form fluxes through 3–cycles of the CY space must fix all its complex structure moduli plus the type IIB dilaton field at string tree level [7]. In the effective action non-vanishing 3–form fluxes generate a tree level N=1 superpotential [8] for the complex structure moduli chiral multiplets and the dilaton, and one has to check that in the corresponding scalar potential the supersymmetry conditions (vanishing of all F–terms) stabilizes all these scalar fields. As it turns out, this condition is generically satisfied in type IIB orientifold compactifications. Without any further input the corresponding supersymmetric flux ground states are flat Minkowski vacua where all the Kähler moduli are still undetermined flat directions of the effective scalar potential.

(ii) In order to stabilize the Kähler moduli of the underlying Calabi-Yau space, non-perturbative effects are needed. The Kähler moduli correspond to the volumes of homologically non-trivial four–cycles inside the Calabi-Yau manifold, called divisors. Specifically, a non-vanishing superpotential for these fields may be generated by Euclidean $D3$–brane instantons which are wrapped around the corresponding divisors [9]. For the divisor volume to be stabilized by Euclidean $D3$–instantons, it must fulfill a certain condition: it must allow for precisely two fermionic zero modes on the $D3$–brane world volume. As we will show, these conditions cannot be satisfied for all orientifolds of blown–up orbifolds geometries, and hence we are left with models, which still possess flat directions in the Kähler moduli space. Another, alternative mechanism to stabilize Kähler moduli fields is provided by gaugino condensation [10], within the effective Yang-Mills gauge theory on a stack of space-time filling $D7$–branes, which are also wrapped around a divisor of the CY space. Again certain conditions have to be met for a non–vanishing gaugino condensate. First, being space-time filling, the $D7$–branes have to satisfy together with the $O7$–orientifold planes the RR–tadpole conditions. This restricts the number of $D7$–branes and also the choice of divisors around which they can be wrapped. Second, the matter spectrum of the effective $D7$–brane gauge theory must be such that a non-perturbative superpotential is formed by the gaugino condensates. Specifically, no massless adjoint chiral multiplets should appear in the effective gauge theory. This requirement restricts the bosonic zero modes on the $D7$–brane world volume, *i.e.* the $D7$–branes should not have any open string moduli fields, namely either Wilson line moduli on its world volume or geometrical moduli that determine the $D7$–brane positions in the two-dimensional transversal directions inside the CY space. These conditions essentially mean that the divisor must be rigid inside the CY space. Furthermore also the spectrum of fundamental chiral fields is restricted, such as the number of bifundamental representations that originate from $D7$–brane intersections or from open string 2–form fluxes on the $D7$–brane world volumes.

(iii) Even if one has succeeded to stabilize all moduli in an AdS vacuum, there is another condition that is related to the stability of the orientifold compactification after the uplift to a dS vacuum. Concretely, it is well known that stable AdS vacua are also possible for the case that the $(\text{mass})^2$ matrix M_{ij}^2 of the moduli contains negative eigenvalues as long as the absolute value of M_{ij}^2 is within the Breitenlohner–Freedman bound [11]. However if one wants to perform the dS uplift in the way described above, we have to insist that M_{ij}^2 is positive definite. As we will discuss, this requirement gives rise to a quite powerful theorem: for CY orientifold compactifications without complex structure moduli one can show that under very mild conditions on the Kähler potential of the Kähler moduli fields, stability after the uplift cannot be achieved.

In order to see for which resolved orbifold geometries all moduli can be stabilized we will essentially follow the following steps:

- We start with some toroidal orbifold and describe the local patches near the singularity with toric methods. Then we resolve the singularities via blow-ups. Next we have to glue together the local patches in order to get a smooth CY space Y_6. Having done this we can compute certain geometrical and topological data of Y_6. These are the triple intersection numbers of the divisors, which enter the Kähler potential of the Kähler moduli fields. In addition one also needs the topological Hodge numbers of the divisors, in order to decide which divisors can contribute to the non-perturbative superpotential. This first step is subject of the companion paper [4] on the geometry of blown-up orbifold spaces. In Section 5 we will heavily rely on these results.

- In the second step we perform an orientifold projection on the smooth CY, which gives rise to $O3/O7$–planes and also $D3$– and $D7$–branes for tadpole cancellation. Going from N=2 type IIB compactifications on a CY covering space Y_6, such as the blown up orbifolds, to its N=1 relative on a CY orientifold X_6 with $O3$– and $O7$–planes leads to several new effects which are relevant in the context of the KKLT moduli stabilization program. First of all, the orientifold projection leads to a field redefinition of the Kähler moduli fields together with the dilaton field. Second, the $\mathbf{Z}_2$ orientifold projection will change some of the triple intersection numbers, which enter the Kähler potential of the Käher moduli fields. Third, whenever an orbifold group contains subgroups of odd order, some of the fixed sets under these subgroups will not be invariant under the global orientifold $\mathbf{Z}_2$ involution. *E.g.* when certain fixed points or lines are identified under the orientifold quotient, the 2–(co)homology splits into an invariant and an anti-invariant part under the orientifold action. This implies that only the geometric divisor moduli of invariant patches can be fixed by the non-perturbative superpotential. On the other hand, the moduli associated to non-invariant divisors are not any more geometrical divisor volumes, but rather are given by a combination of two-form gauge potentials. Since the superpotential does not depend on them, these moduli cannot be fixed by the KKLT mechanism. However, as we will discuss, they will appear in a D–term scalar potential. Requiring supersymmetric ground-states with vanishing D–terms will allow us to stabilize also these moduli fields.

- Then we determine the Kähler potential for the Kähler moduli fields using the triple intersection numbers. In addition one has to compute also the Kähler potential for the complex structure moduli fields. We will be able to do this explicitly only for orbifolds with only untwisted complex structure moduli fields.

- Furthermore we compute the non-perturbative superpotential for the Kähler moduli fields from the divisor topologies. In addition we compute the tree level 3–form flux superpotential for the dilaton and the complex structure moduli fields.

- Finally we determine the supersymmetric AdS vacua of the scalar potential and compute the corresponding values of all moduli fields.

2. Moduli stabilization and stability in type *IIB* orientifolds

2.1. Calabi–Yau orientifolds of type IIB with D3/D7–branes

We start with a type IIB compactification on a Calabi–Yau (CY) manifold Y_6. This leads to N=2 supersymmetry in $D=4$ dimensions. The geometry of the manifold Y_6 is described by $h_{(1,1)}(Y_6)$ Kähler moduli and $h_{(2,1)}(Y_6)$ complex structure moduli. These moduli fields represent scalar components of N=2 hyper– and vector multiplets, respectively. Together with the universal hypermultiplet we have $h_{(1,1)}(Y_6)+1$ hypermultiplets and $h_{(2,1)}(Y_6)$ vector multiplets.

To arrive at N=1 supersymmetry in $D=4$ we introduce an orientifold projection $\mathcal{O}$, which produces orientifold $O3$– and $O7$–planes. To cancel tadpoles and to construct models of phenomenological interest we add $D3$– and $D7$–branes. The orientifold projection $\mathcal{O}$ [12,13]

$$\mathcal{O}=(-1)^{F_L}\ \Omega\ \sigma^\star \tag{2.1}$$

acting on the closed type IIB string states is given by a combination of world–sheet parity transformation Ω and a reflection σ in the internal CY space. The CY geometry Y_6 modded out by the additional involution σ is labeled by X_6. To obtain $O3/O7$–planes the action σ must act holomorphically and satisfy

$$\sigma^\star\ \Omega_{(3,0)}=-\Omega_{(3,0)}\ , \tag{2.2}$$

with $\Omega_{(3,0)}$ the holomorphic 3–form of the CYM X_6.

Due to the holomorphic action of σ, the latter splits the cohomology groups $H^{(p,q)}(Y_6)$ into a direct sum of an even eigenspace $H_+^{(p,q)}(X_6)$ and an odd eigenspace $H_-^{(p,q)}(X_6)$ [13]. Since the Kähler form J is invariant under the orientifold action, it is expanded w.r.t. a basis of $H_+^{(1,1)}(X_6)$. On the other hand, because of (2.2), the holomorphic 3–form $\Omega_{(3,0)}$ may be expanded w.r.t. a real symplectic basis $(\alpha_\lambda,\beta^\lambda)$ of $H_-^{(3)}(X_6)$, *i.e.*

$$J=\sum_{k=1}^{h_{(1,1)}^{(+)}(X_6)} t^k\ \omega_k\quad,\quad \Omega_{(3,0)}=\sum_{\lambda=0}^{h_{(2,1)}^{(-)}(X_6)} X^\lambda\alpha_\lambda-F_\lambda\beta^\lambda\ , \tag{2.3}$$

with (X^λ,F_λ) the periods of the original CYM Y_6. Furthermore, in type IIB orientifolds with $D3$– and $D7$–branes, the $NSNS$ two–form B_2 and the RR two–form C_2 are odd under the orientifold action $(-1)^{F_L}\Omega$. Hence, they are expanded w.r.t. a basis of the cohomology $H_-^{(1,1)}(X_6)$, *i.e.*

$$B_2=\sum_{a=1}^{h_{(1,1)}^{(-)}(X_6)} b^a\ \omega_a\quad,\quad C_2=\sum_{a=1}^{h_{(1,1)}^{(-)}(X_6)} c^a\ \omega_a\ . \tag{2.4}$$

In type IIB orientifolds the fields b^a and c^a give rise to $h^{(-)}_{(1,1)}(X_6)$ complex scalars

$$G^a = i\ c^a - S\ b^a \quad , \quad a = 1, \ldots, h^{(-)}_{(1,1)}(X_6) \tag{2.5}$$

of N=1 chiral multiplets [14], whose vevs eventually should be fixed. Clearly, $D3$– and $D7$–branes may be wrapped only around 4–cycles whose Poincaré dual 2–form is an element of $H^2_+(X_6)$. To summarize, in addition to the dilaton field

$$S = e^{-\phi_{10}} + i\ C_0 \tag{2.6}$$

a CY orientifold compactification X_6 has $h^{(+)}_{(1,1)}(X_6)$ Kähler moduli t^k, $h^{(-)}_{(1,1)}(X_6)$ scalars G^a and $h^-_{(2,1)}(X_6)$ complex structure moduli u^λ. As shown[1] in Table 1, under the orientifold action $\mathcal{O}$ the original set of $h_{(1,1)}(Y_6) + 1$ N=2 hypermultiplets and $h_{(2,1)}(Y_6)$ N=2 vectormultiplets is split into a set of N=1 chiral and vectormultiplets.

1	dilaton	S	chiral multiplet	$\int_{X_6} \Omega \wedge G_3$	$ISD\ 3{-}form\ flux\ G_3$
$h^{(-)}_{(2,1)}(X_6)$	CS moduli	u^λ	chiral multiplets	$\int_{X_6} \Omega \wedge G_3$	$ISD\ 3{-}form\ flux\ G_3$
$h^{(+)}_{(1,1)}(X_6)$	Kähler moduli	t^k, ρ^k	chiral/linear multiplets	e^{-T}	$D3$ instanton gaugino condensation
$h^{(-)}_{(1,1)}(X_6)$	add. moduli	b^a, c^a	chiral multiplets	$\int_{C_4} J \wedge B_2$ $(D_\mu G^a)^2$	calibration massive vector
$h^{(+)}_{(2,1)}(X_6)$	add. vectors	$\widetilde{V}^j_\mu$	vector multiplets	–	–

Table 1: *Moduli of Calabi–Yau orientifold X_6 and their stabilization mechanism.*

The additional $h^{(+)}_{(2,1)}(X_6)$ vectors (and their magnetic duals) arise from the Ramond 4–form C_4 reduced w.r.t. the cohomology $H^{(3)}_+(X_6)$. Besides the dilaton field S in the Kähler potential for the moduli fields

$$K = -\ln(S + \overline{S}) - 2\ln\left(\frac{1}{6}\int_{X_6} J \wedge J \wedge J\right) - \ln\left(-i\int_{X_6} \Omega \wedge \overline{\Omega}\right) \tag{2.7}$$

only the $h^{(+)}_{(1,1)}(X_6)$ invariant Kähler moduli t^k and the $h^-_{(2,1)}(X_6)$ invariant complex structure moduli enter explicitly. However, the string theoretical Kähler moduli t^j are not yet scalars of an N=1 chiral multiplet. After defining the proper holomorphic moduli fields T^j (in the string frame[2]) [14]

---

[1] Here and in the following, where no confusion with the orientifold action Ω may occur we shall use Ω for $\Omega_{(3,0)}$.

$$T^j = \frac{3}{4}\ \mathcal{K}_{jkl}\ t^k\ t^l - \frac{3}{8}\ e^{\phi_{10}}\ \mathcal{K}_{jbc}\ \overline{G}^b\ (G+\overline{G})^c + \frac{3}{2}\ i\ \left(\ \rho^j - \frac{1}{2}\ \mathcal{K}_{jbc}\ c^b\ b^c\ \right)\ , \tag{2.8}$$

the second term $K_{KM} = -2\ln Vol(X_6) = -2\ln\frac{1}{6}\mathcal{K}_{ijk}t^it^jt^k$ in (2.7) may be expressed in terms of the N=1 fields T^j. This way, in the low–energy effective action of type IIB CY orientifolds, the fields G^a do enter the Kähler potential for the Kähler moduli t^k through eliminating the moduli t^k via the definition (2.8). By that the Kähler potential K_{KM} for the $h^{(+)}_{(1,1)}(X_6)$ Kähler moduli T^j becomes a complicated function $K_{KM}(S,T^j,G^a)$ depending on the dilaton S, the $h^{(+)}_{(1,1)}(X_6)$ moduli T^j and the $h^{(-)}_{(1,1)}(X_6)$ moduli G^a [14]. In (2.8) the axion ρ^j originates from integrating the RR 4–form along the 4–cycle C_j. The full Kähler potential

$$K = -\ln(S+\overline{S}) - 2\ln Vol(X_6) + K_{CS} \tag{2.9}$$

for the dilaton S, $h^{(+)}_{(1,1)}(X_6)$ Kähler moduli T^k, $h^{(-)}_{(1,1)}(X_6)$ scalars G^a and $h^{(-)}_{(2,1)}(X_6)$ complex structure moduli takes the form [14]:

$$K = -\ln(S+\overline{S}) + K_{KM}(S,T^j,G^a) + K_{CS}\ . \tag{2.10}$$

To illustrate the structure of the modified Kähler potential, let us briefly discuss the case $h^{(+)}_{(1,1)}(X_6) = 1 = h^{(-)}_{(1,1)}(X_6)$. The Kähler potential for the single Kähler modulus t is: $K_{KM}(t) = -2\ln t^3$. With the intersection numbers $\mathcal{K}_{ttt} = 6$, $\mathcal{K}_t = 6t^2$ and $\mathcal{K}_{tbb} = 1$ we derive from (2.8)

$$T = \frac{9}{2}\ t^2 - \frac{3}{8}\ e^{\phi_{10}}\ \overline{G}\ (G+\overline{G}) + \frac{3}{2}\ i\ \left(\rho - \frac{1}{2}\ c\ b\right)\ ,$$

and the full Kähler potential (2.10) becomes:

$$K = -\ln(S+\overline{S}) - 3\ \ln\frac{1}{9}\left[\ T+\overline{T}+\frac{3}{4}\ \frac{(G+\overline{G})^2}{S+\overline{S}}\ \right] + K_{CS}\ . \tag{2.11}$$

Before adding background fluxes, in the effective $D=4$ action the fields S, u^λ, t^j, b^a and c^a have flat directions, *i.e.* no potential is generated for them and their vevs may assume arbitrary values in their moduli spaces. Fixing these moduli through some $F-$ or D–term potential is the main topic of this article. In the two last columns of Table 1 we have shown the different mechanisms how to stabilize these moduli. These mechanisms shall be discussed in Subsections 2.2, 2.3 and Section 3.

---

[2] In the Einstein frame the Kähler moduli t^k are multiplied with $e^{-\frac{1}{2}\phi_{10}}$. In the Einstein frame the CY volume reads $Vol(X_6) = \frac{1}{6}e^{-\frac{3}{2}\phi_{10}}\ \mathcal{K}_{ijk}t^it^jt^k$.

2.2. Stabilization of Kähler moduli associated to the cohomology $H^{(1,1)}_+(X_6)$

In the following we shall assume a flux compactification of a type IIB CY orientifold with $h^{(-)}_{(1,1)}(X_6) = 0$ and a general Kähler potential K_{KM} for the $n := h^{(+)}_{(1,1)}(X_6)$ Kähler moduli:

$$K_{KM}(T^1,\ldots,T^n,\overline{T}^1,\ldots,\overline{T}^n) = K(T^1+\overline{T}^1,\ldots,T^n+\overline{T}^n)\ . \tag{2.12}$$

We shall adress the case $h^{(-)}_{(1,1)}(X_6) \neq 0$ in Section 3. We consider the racetrack superpotential [15]:

$$W = W_0(S,U) + \lambda\ \sum_{j=1}^{n} \gamma_j(S,U)\ e^{a_j\ T^j}\ . \tag{2.13}$$

The first term W_0 of (2.13) represents the tree–level flux superpotential [8]

$$W_0(S,U) = \int_{X_6} G_3 \wedge \Omega \tag{2.14}$$

depending on the dilaton S and the $h^{(-)}_{(2,1)}(X_6)$ complex structure moduli U^λ. Since $\Omega \in H^{(3)}_-(X_6)$, we also must have $G_3 \in H^{(3)}_-(X_6)$ with $2h^{(-)}_{2,1}(X_6)+2$ complex flux components. In *Eq.* (2.13) we assume $W_0 \in \mathbf{C}$, $\gamma_j \in \mathbf{C}$, and $a_j \in \mathbf{R}_-$. In addition, $\lambda \in \mathbf{R}$ is a real parameter accounting for a possible so–called Kähler gauge (*cf.* Subsection 5.4). The latter may be used to adjust a certain flux value given by W_0 to a given minimum in the Kähler moduli space $(T^1,\ldots,T^n)$, *cf. Ref.* [16]. We do not consider a possible open string moduli dependence of the superpotential [17,18].

The work of KKLT [1] proposes a mechanism to stabilize all moduli at a small positive cosmological constant. This procedure is accomplished through three steps. One first dynamically fixes the dilaton S and the complex structure moduli U^λ through the tree–level piece W_0 (given in *Eq.* (2.14)) of the superpotential. This is accomplished with a generic 3–form flux G_3 with both ISD– and $IASD$–flux components. At the minimum of the scalar potential in the complex structure and dilaton directions, the flux becomes ISD and the potential assumes the value $V_0(S,U^\lambda) = -3e^K|W_0|^2$. The soft masses m_S, m_{U^λ} for the dilaton and complex structure scalars are generically of the order α'/R^3 [19]. In the large radius approximation $\mathrm{Re}(T) \gg 1$, the non–perturbative terms in (2.13) only amount to a small exponentially suppressed additional contribution to m_S, m_{U^λ}. According to [20] the latter is negligible. The second step is the addition of the non–perturbative piece to the superpotential (2.13), which allows the stabilization of the Kähler moduli T^j at a supersymmetric AdS minimum. The soft masses for the Kähler moduli are much smaller than soft masses m_S and m_{U^λ}. This property allows us to separate the first and second step, *i.e.* to effectively first integrate out the dilaton and complex structure moduli. Nonetheless, strictly speaking these two steps should be treated at the same time and that is what we shall do in Subsection 2.3. The stability of AdS vacua in gravity coupled to scalar fields has been investigated in *Ref.* [11]. Stability is guaranteed, if all scalar masses fulfill the Breitenlohner–Freedman (BF) bound [11], *i.e.* their mass eigenvalues do not fall below a certain minimal bound. The latter is a negative number related to the scalar potential at the minimum. It can be shown in a completely model independent way that all scalars have masses above this bound at any N=1 supersymmetric AdS minimum in supergravity theories (*cf. e.g.* [21] and Appendix C of *Ref.* [22]). However, the third and final step in the KKLT scenario consists in the addition of one anti $D3$–brane, *i.e.* a positive contribution to the scalar potential, which lifts the AdS minimum to a dS minimum. The masses for the moduli fields do not change significantly during this process. However stable dS vacua require positive mass eigenvalues. Hence, any negative mass eigenvalue before the uplift is unacceptable since the effect of the anti $D3$–branes on the mass eigenvalues is too small to change a negative mass to positive.

In (2.13), the sum of exponentials accounts for $D3$–brane instantons and gaugino condensation on stacks of $D7$–branes. The $D3$–instantons come from wrapping (Euclidean) $D3$–branes on internal 4–cycles C_4^j of the CY orientifold X_6. The latter have the volume $\mathrm{Re}(T^j)$ and lead to the instanton effect $e^{-2\pi T^j}$ in the superpotential, *i.e.* $a_j = -2\pi$. The gauge coupling on a $D7$–brane which is wrapped on the 4–cycle C_4^j is given by $\mathrm{Re}(T^j)$, *cf. Eq.* (2.8). Hence, gaugino condensation on this $D7$–brane yields the effect e^{-T^j/b_a} in the superpotential. *E.g.* for the gauge group $SU(M)$ we have $b_{SU(M)} = \frac{M}{2\pi}$, *i.e.* $a_j = -\frac{2\pi}{M}$. On the $D7$–brane, $\gamma_j(S,U)$ may comprise one–loop effects and further instanton effects from $D(-1)$–branes: One loop corrections to the gauge coupling give rise to [23]

$$\gamma_j \sim \eta(U^\lambda)^{-2/b_a} \ , \tag{2.15}$$

while additional instantons in the $D7$–gauge theory amount to:

$$\gamma_j \sim e^{-S/b_a \int_{C_4^j} F\wedge F} \ . \tag{2.16}$$

Supersymmetric vacuum solutions are found by finding the zeros of the F–terms: $\overline{F}^{\overline{M}} = K^{\overline{M}J}\ (\partial_J W + W\ K_J)$. Solutions to the equations $F^M = 0$ give rise to extremal points of the scalar potential. In addition, it has to be verified whether those zeros lead to a stable minimum. Since the matrix $K^{\overline{M}J}$ is positive definite, the zeros $(T_0^1, \ldots, T_0^n)$ in the Kähler moduli space are determined by the n equations:

$$\partial_{T^j} W + W\ K_{T^j} = 0 \quad , \quad j = 1, \ldots, n \tag{2.17}$$

following from the requirement of vanishing F–terms. These equations turn into the $h^{(+)}_{(1,1)}(X_6)$ relations

$$\lambda\,\gamma_i=-\left(\prod_{j\neq i}^{n}|a_j|\right)\ e^{-a_i\ T^i}\ W_0\ \left.\frac{K_{T^i}}{\sum\limits_{j=1}^{n}K_{T^j}\ \prod\limits_{\substack{k=1\\k\neq j}}^{n}|a_k|-\prod\limits_{k=1}^{n}|a_k|}\right|_{T^l=T^l_0}\quad ,\quad i=1,\ldots,n \tag{2.18}$$

to be satisfied at this extremum. Since K_{T^i} and a_i are real, from *Eq.* (2.18) we may easily deduce the vevs t^j_2 of the axions at the AdS–minimum. After introducing the phases $W_0=|W_0|\ e^{i\varphi}$and $\gamma_i=|\gamma_i|\ e^{i\phi_i}$ we obtain $(T^j=t^j_1+it^j_2)$

$$t^i_2=\frac{1}{a_i}\ \left[\ \varphi+\pi\ (1+\rho^i)-\phi_i\ \right]+\frac{2\pi}{a_i}\ \mathbf{Z}\quad ,\quad i=1,\ldots,n \tag{2.19}$$

as vevs for the axion fields. Above we have introduced the numbers

$$\rho^i=\frac{1}{\pi}\ arg\left(\frac{\lambda\ K_{T^i}}{\sum\limits_{j=1}^{n}K_{T^j}\ \prod\limits_{\substack{k=1\\k\neq j}}^{n}|a_k|-\prod\limits_{k=1}^{n}|a_k|}\right)\in\{0,1\}\ .$$

For the case that an exponential $e^{a_jT^j}$ accounts for gaugino condensation in an $SU(M)$ gauge group, we have $a_j=-\frac{2\pi}{M}$ and in *Eq.* (2.19) the additional shift $\frac{2\pi}{a_j}$ $\mathbf{Z}$ becomes M $\mathbf{Z}$. The latter becomes trivial, if the Kähler modulus T^j enjoys a discrete shift symmetry, *e.g.* $T^j\rightarrow T^j+1$. On the other hand, if no such symmetry exists, in *Eq.* (2.19) the additional shifts $\frac{2\pi}{a_j}$ $\mathbf{Z}$ give rise to an infinite number of extrema obtained from one another by shifts in the axionic directions t^j_2. A useful relation to be satisfied at the extremum is the ratio

$$\frac{\gamma_i}{\gamma_j}=e^{a_j\ T^j-a_i\ T^i}\ \frac{a_j}{a_i}\ \frac{K_{T^i}}{K_{T^j}}\quad ,\quad i,j=1,\ldots,n\ . \tag{2.20}$$

The latter equation may be written as

$$\frac{|\gamma_i|}{|\gamma_j|}=e^{i\phi_{ji}}\ e^{a_j\ t^j_1-a_i\ t^i_1}\ \frac{a_2}{a_1}\ \frac{K_{T^1}}{K_{T^2}}\quad ,\quad \phi_{ji}=\phi_j-\phi_i+a_j\ t^j_2-a_i\ t^i_2\ . \tag{2.21}$$

Since $\phi_{ji}\in\{0,\pi\}$, the directions of the axions t^j_2 strongly depend on the signs of the first Kähler derivatives K_{T^j} and the phases ϕ_j of the coefficients γ_j.

Moreover, from the relation (2.19) we see that any complex phase of W_0 and γ_i may be absorbed into a redefinition of the axion vev at the minimum. Hence, in the following we may assume without any restriction:

$$W_0\in\mathbf{R}^+\quad ,\quad \gamma_j\in\mathbf{R}^+\ .$$

Finally, at the extremum $(T^1_0,\ldots,T^n_0)$, the scalar potential $\widetilde{V}(T^1,\overline{T}^1,\ldots,T^n,\overline{T}^n)$ assumes the negative value

$$\widetilde{V}_{min} = -3\ \ e^K \left(\prod_{k=1}^{h^{(+)}_{(1,1)}(X_6)} \ a_k^2 \right) \frac{|W_0|^2}{\left(\sum\limits_{j=1}^{n} K_{T^j} \prod\limits_{\substack{k=1\\k\neq j}}^{n} |a_k| - \prod\limits_{k=1}^{n} |a_k| \right)^2} \Bigg|_{T^l=T^l_0} \tag{2.22}$$

2.3. (In)stability of flux compactifications without complex structure moduli

In this subsection we shall continue and generalize the discussion initiated by the research [24,3] (see also *Refs.* [25,26]) on the stability criteria in a KKLT setup [1]. In these references, the conditions under which a stable vacuum may be found have been investigated for some selected orbifold examples. This means in particular that a special Kähler potential for the underlying coset spaces parametrizing the Kähler moduli space has been used. The general property for those examples is that without any complex structure modulus or with fixed complex structures, as it is the case for many toroidal orbifold examples at the orbifold point, no stable vacuum in a KKLT setup with a non–perturbative superpotential exists. The question we shall address here is, whether this is a general feature, independent of the form of the Kähler potential of compactifications without complex structure moduli, in particular also holding for the resolved orbifolds. Note, that in those cases, the Kähler potential assumes a quite different form than for the coset spaces.

For the case of $h^{(-)}_{(2,1)}(X_6) = 0$ and $h^{(+)}_{(1,1)}(X_6) =: n$ the superpotential (2.13) assumes the form:

$$W = B + A\ S + \lambda \sum_{i=1}^{h^{(+)}_{(1,1)}(X_6)} \ \gamma_i\ e^{a_i\ T^i}\ . \tag{2.23}$$

Generically, we have $A, B \in \mathbf{C}$, $\gamma_j \in \mathbf{C}$, and $a_i \in \mathbf{R}_-$. However, we have argued in the previous subsection that without any restriction we might assume $A, B, \gamma_j \in \mathbf{R}^+$. Nevertheless, we shall treat them in the following as arbitrary complex parameters. In addition, $\lambda \in \mathbf{R}$ is a real parameter accounting for a possible so–called Kähler gauge, as discussed in [16].

The supersymmetric vacuum solutions are given by solving the equations $F^M = 0$ for $M = S, T^1, \ldots, T^n$. With the Kähler potential

$$-\ln(S+\overline{S}) + K_{KM}(T^1,\ldots,T^n,\overline{T}^1,\ldots,\overline{T}^n)$$

for the moduli space $(S, T^1, \ldots, T^n)$ we obtain the F–term

$$\overline{F}_S = e^{\frac{1}{2}K} \; (S+\overline{S})^{1/2} \; \left(-B + A\;\overline{S} - \lambda \sum_{i=1}^{h^{(+)}_{(1,1)}(X_6)} \gamma_i \; e^{a_i\;T^i} \right) \tag{2.24}$$

for the dilaton field S. The solutions $(S_0, T^1_0, \ldots, T^n_0)$ to the vacuum equations

$$F^M = 0 \quad , \quad M = S, T^1, \ldots, T^n \tag{2.25}$$

are given by the dilaton vev S_0

$$S_0 = -\frac{\prod\limits_{k=1}^{h^{(+)}_{(1,1)}(X_6)} |a_i|}{2\;|A|^2} \; \frac{\overline{A}\;B + A\;\overline{B}}{2 \sum\limits_{j=1}^{h^{(+)}_{(1,1)}(X_6)} K_{T^j} \prod\limits_{\substack{k=1\\ k\neq j}}^{h^{(+)}_{(1,1)}(X_6)} |a_k| - \prod\limits_{k=1}^{h^{(+)}_{(1,1)}(X_6)} |a_k|} - \frac{1}{2\;|A|^2}\;(\overline{A}\;B - A\;\overline{B}) \; , \tag{2.26}$$

and the n relations:

$$\begin{aligned} \lambda\;\gamma_i &= -\frac{\prod\limits_{j\neq i}^{h^{(+)}_{(1,1)}(X_6)} |a_j|}{\overline{A}} \; e^{-a_i\;T^i} \; \frac{K_{T^i}\;(\overline{A}\;B + A\;\overline{B})}{2 \sum\limits_{j=1}^{h^{(+)}_{(1,1)}(X_6)} K_{T^j} \prod\limits_{\substack{k=1\\ k\neq j}}^{h^{(+)}_{(1,1)}(X_6)} |a_k| - \prod\limits_{k=1}^{h^{(+)}_{(1,1)}(X_6)} |a_k|} \Bigg|_{T^l = T^l_0} \\ &= -\frac{2\;A\;\mathrm{Re}(S)}{a_i}\; e^{-a_i\;T^i}\; K_{T^i} \; \Bigg|_{T^l = T^l_0} \quad , \quad i = 1, \ldots, n\;. \end{aligned} \tag{2.27}$$

Similarly as in the previous subsection (*cf. Eq.* (2.19)) we may determine the vevs for the axion fields. With $A = |A|\;e^{i\varphi}$ and $\gamma_i = |\gamma_i| e^{i\phi_i}$ we obtain:

$$t^i_2 = \frac{1}{a_i}\;[\;\varphi + \pi - \phi_i\;] + \frac{2\pi}{a_i}\;\mathbf{Z} \quad , \quad i = 1, \ldots, n\;. \tag{2.28}$$

Note, that the above expression is somewhat simpler than (2.19), since we have solved for $n+1$ equations $F^M = 0$ thus allowing to eliminate some more terms. In fact, the additional relation, valid at the point $(S_0, T^1_0, \ldots, T^n_0)$, is:

$$\left(-2 \sum_{j=1}^{h^{(+)}_{(1,1)}(X_6)} K_{T^j} \prod_{\substack{k=1\\ k\neq j}}^{h^{(+)}_{(1,1)}(X_6)} |a_k| + \prod_{k=1}^{h^{(+)}_{(1,1)}(X_6)} |a_k| \right) = \frac{(\overline{A}\;B + A\;\overline{B}) \prod\limits_{k=1}^{h^{(+)}_{(1,1)}(X_6)} |a_i|}{2\;|A|^2\;\mathrm{Re}(S)}\;. \tag{2.29}$$

The scalar potential becomes

$$\begin{aligned} V(S,\overline{S},T^1,\overline{T}^1,\ldots,T^n,\overline{T}^n) = {} & \frac{\widetilde{V}(T^1,\overline{T}^1,\ldots,T^n,\overline{T}^n)}{S+\overline{S}} \\ & + \frac{e^K}{S+\overline{S}} \left| B - A\ \overline{S} + \lambda \sum_{k=1}^{h^{(+)}_{(1,1)}(X_6)} \gamma_k\ e^{a_k\ T^k} \right|^2 , \end{aligned} \tag{2.30}$$

with $\widetilde{V}(T^1,\overline{T}^1,\ldots,T^n,\overline{T}^n)$ the scalar potential computed in the Kähler moduli sector $(T^1,\ldots,T^n)$ with Kähler potential K_{KM}. At the vacuum solution $(S_0,T_0^1,\ldots,T_0^n)$, the second piece of (2.30), which is proportional to $|F_S|^2$, vanishes and the scalar potential assumes the value

$$\begin{aligned} V_{min} = \frac{\widetilde{V}_{min}}{S_0+\overline{S}_0} & = -6\ |A|^2\ \mathrm{Re}(S)\ e^K \Big|_{\substack{S=S_0\ , \\ T^l=T_0^l}} \\ & = -3\ \ \frac{e^K}{S+\overline{S}}\ \frac{\prod_{k=1}^{h^{(+)}_{(1,1)}(X_6)} a_k^2}{|A|^2}\ \frac{(\overline{A}\ B + A\ \overline{B})^2}{\left(2 \sum_{j=1}^{h^{(+)}_{(1,1)}(X_6)} K_{T^j} \prod_{\substack{k=1 \\ k\neq j}}^{h^{(+)}_{(1,1)}(X_6)} |a_k| - \prod_{k=1}^{h^{(+)}_{(1,1)}(X_6)} |a_k| \right)^2} \Bigg|_{\substack{S=S_0\ , \\ T^l=T_0^l}} , \end{aligned} \tag{2.31}$$

with $\widetilde{V}_{min}$ given in (2.22).

To analyze the stability of the vacuum solution, we have to calculate the second derivatives of the scalar potential $V(S,\overline{S},T^1,\overline{T}^1,\ldots,T^n,\overline{T}^n)$ at the supersymmetric point. After introducing $S=s_1+is_2$, we find that the mixed derivatives

$$\frac{\partial^2 V}{\partial t_1^i \partial t_2^j} = \frac{\partial^2 V}{\partial t_1^i \partial s_2} = \frac{\partial^2 V}{\partial t_2^i \partial s_1} = \frac{\partial^2 V}{\partial s_1 \partial s_2} = 0 \quad , \quad i,j=1,\ldots,n \tag{2.32}$$

vanish at the extremum $(S_0,T_0^1,\ldots,T_0^n)$. Now we derive the necessary criteria for the matrix encoding the scalar masses to have positive eigenvalues at the point $(S_0,T_0^1,\ldots,T_0^n)$. We have to determine the second derivatives of the potential (2.30) and check, whether the mass matrix is positive definite. In particular, due to (2.32), the mass matrix M

$$M = \begin{pmatrix} \frac{\partial^2 V}{\partial (t_1^{i_1})^2} & \cdots & \frac{\partial^2 V}{\partial t_1^{i_1} \partial t_1^n} & \frac{\partial^2 V}{\partial t_1^n \partial s_1} \\ \vdots & \vdots & \vdots & \vdots \\ \frac{\partial^2 V}{\partial t_1^n \partial t_1^{i_1}} & \cdots & \frac{\partial^2 V}{\partial (t_1^n)^2} & \frac{\partial^2 V}{\partial t_1^n \partial s_1} \\ \frac{\partial^2 V}{\partial s_1 \partial t_1^{i_1}} & \cdots & \frac{\partial^2 V}{\partial s_1 \partial t_1^n} & \frac{\partial^2 V}{\partial s_1^2} \end{pmatrix} \tag{2.33}$$

of the axions $s_2, t_2^1, \ldots, t_2^n$ has to be positive definite. For the moment we shall concentrate on the properties of this axionic mass matrix M. We shall derive a general expression for the determinant $\det(M)$ of the axion mass matrix M and express it in terms of its principal minors. To tackle this project, we first introduce the 2^n-2 principal submatrices of M:

$$M_{i_1\ldots i_p} = \begin{pmatrix} \frac{\partial^2 V}{\partial (t_1^{i_1})^2} & \cdots & \frac{\partial^2 V}{\partial t_1^{i_1}\partial t_1^{i_p}} & \frac{\partial^2 V}{\partial t_1^{i_1}\partial s_1} \\ \vdots & \vdots & \vdots & \vdots \\ \frac{\partial^2 V}{\partial t_1^{i_p}\partial t_1^{i_1}} & \cdots & \frac{\partial^2 V}{\partial (t_1^{i_p})^2} & \frac{\partial^2 V}{\partial t_1^{i_p}\partial s_1} \\ \frac{\partial^2 V}{\partial s_1\partial t_1^{i_1}} & \cdots & \frac{\partial^2 V}{\partial s_1\partial t_1^{i_p}} & \frac{\partial^2 V}{\partial s_1^2} \end{pmatrix} \quad , \quad \begin{array}{c} p=1,\ldots,n-1 \ , \\ 1 \leq i_1 < \ldots < i_p \leq n \end{array} \ . \tag{2.34}$$

Any principal submatrix of a positive definite matrix must be positive definite [27]. Therefore, all the submatrices $M_{i_1}, M_{i_1 i_2}, \ldots, M_{i_1\ldots i_{n-1}}$ must have positive determinant, *i.e.* those principal minors of M must satisfy

$$\det(M_{i_1\ldots i_p}) > 0$$

in order that M may be positive definite. After some tedious algebra the determinant of the matrix M may be expressed[3] in terms of its 2^n-2 submatrices $M_i, M_{ij}, M_{ijk}, \ldots$:

$$\begin{aligned} \det(M) = & - \left[\ \mathrm{Re}(S)\ |A|^2\ e^K\ \right]^n \left(\prod_{j=1}^n a_j\ K_{T^j} \right) \\ & \times \left\{ \ (3\cdot 4)^n\ \frac{\partial^2 V}{\partial s_1^2} + 2\cdot 4^n\ \mathrm{Re}(S)^{-1}\ |A|^2 \left(\prod_{j=1}^n a_j\ K_{T^j} \right) \frac{e^K}{\det(\mathcal{G})} \right. \\ & \left. + \sum_{s=1}^{n-1} (3\cdot 4)^{n-s}\ \left[\ \mathrm{Re}(S)\ |A|^2\ e^K\ \right]^{-s} \sum_{i_1<\ldots<i_s}^{n} \frac{\det(M_{i_1\ldots i_s})}{\prod_{r=1}^{s} a_{i_r}\ K_{T^{i_r}}} \right\} \quad , \quad n \geq 1 \ . \end{aligned} \tag{2.35}$$

Since $a_i < 0$ and $\det(\mathcal{G}) > 0$ from this relation we conclude that the axionic mass matrix M fails to be positive definite, if

$$\boxed{\forall\ \ i = 1, \ldots, h_{(1,1)}^{(+)}(X_6): \qquad K_{T^i} < 0} \tag{2.36}$$

since in that case *Eq.* (2.35) yields $\det(M) < 0$. Hence, no stable uplift is possible, if (2.36) is fulfilled.

---

[3] We introduce the $n \times n$ matrix $\mathcal{G} = \left\{ \partial_{T^i} \partial_{\overline{T}^j} K \right\}_{i,j,=1,\ldots,n}$, which is positive definite.

We may also derive a compact expression for the minors $\det(M_i)$ themselves. For that, we introduce the $n \times n$ matrix

$$\mathcal{O} = \left\{ -K_{T^i \overline{T}^j} + \frac{1}{2} K_{T^i} K_{\overline{T}^j} \right\}_{i,j=1,\ldots,n} \tag{2.37}$$

and its n principal $n-1 \times n-1$ submatrices

$$\mathcal{O}_k = \left\{ -K_{T^i \overline{T}^j} + \frac{1}{2} K_{T^i} K_{\overline{T}^j} \right\}_{\substack{i,j=1,\ldots,n \\ i,j \neq k}} \quad , \quad k = 1, \ldots, h_{(1,1)}^{(+)}(X_6)\ . \tag{2.38}$$

Again, after some tedious work we arrive at the following relation:

$$\begin{aligned} \det(M_i) = &-12\ |A|^2\ \mathrm{Re}(S)\ e^K\ a_i\ K_{T^i}\ \frac{\partial^2 V}{\partial s_1^2} \\ &+ 8\ |A|^4\ e^{2K}\ a_i^2\ K_{T^i}^2\ \frac{(-1)^n\ \det(\mathcal{O}_i)}{\det(\mathcal{G})} \quad , \quad i = 1, \ldots, h_{(1,1)}^{(+)}(X_6)\ . \end{aligned} \tag{2.39}$$

Since $a_i < 0$ and $\det(\mathcal{G}) > 0$ from this equation we conclude: If

$$\boxed{\exists\ \ i \in \{\ 1, \ldots, h_{(1,1)}^{(+)}(X_6)\ \} : \qquad K_{T^i} < 0 \quad \bigwedge\ \ (-1)^n\ \det(\mathcal{O}_i) \leq 0} \tag{2.40}$$

holds, no stable uplift is possible, either.

Note, that the two (independent) criteria (2.36) and (2.40) do not depend on the flux parameters A and B and the complex coefficients γ_j in (2.23). The Kähler potential (2.12) is enough to check the stability criteria of the KKLT setup (2.23). According to (2.36) and (2.40) the condition of a positive definite axion mass matrix boils down to properties of the first derivatives K_{T^j} at the AdS vacuum solution (2.25). The conditions (2.36) and (2.40) are two *independent* criteria for the AdS solution $(S_0, T_0^1, \ldots, T_0^n)$ *not* yielding positive masses for the axions. However, if the conditions are not met, for a KKLT scenario there are further criteria to be checked. The latter arise from calculating the masses for the real parts of $S, T^1, \ldots, T^n$ and verifying their positivity. To summarize, for a KKLT setup with n Kähler moduli and no complex structure modulus, a controlled uplift is *not possible* if at least one of the two conditions (2.36) and (2.40) on the Kähler potential (2.12) is met.

For CY orientifolds X_6 with N=1 supersymmetry we may rewrite the inequalities (2.36) such, that they become conditions on the original Calabi–Yau moduli t^j relevant for type *IIB* compactifications without orientifold projection. As discussed in Subsection 2.1, the orientifold (2.1) splits the forms ω_i into $h_{(1,1)}^{(+)}(X_6)$ elements ω_i of $H_+^{(1,1)}(X_6)$ and

$h^{(-)}_{(1,1)}(X_6)$ elements of $H^{(1,1)}_-(X_6)$. The correct Kähler moduli T^j arise through a Legendre transform on the original Calabi–Yau volume $Vol(t^i)$ [28]: With the Kähler form

$$J = \sum_{j=1}^{h^{(+)}_{(1,1)}(X_6)} t^j \ \omega_j \tag{2.41}$$

the following intersection numbers may be introduced:

$$\begin{aligned} \mathcal{K} &= \int_{X_6} J \wedge J \wedge J = \mathcal{K}_{ijk}\ t^i\ t^j\ t^k\ ,\ \mathcal{K}_i = \int_{X_6} \omega_i \wedge J \wedge J = \mathcal{K}_{ijk}\ t^j\ t^k, \\ \mathcal{K}_{ij} &= \int_{X_6} \omega_i \wedge \omega_j \wedge J = \mathcal{K}_{ijk}\ t^k \quad , \quad \mathcal{K}_{ijk} = \int_{X_6} \omega_i \wedge \omega_j \wedge \omega_k\ , \end{aligned} \tag{2.42}$$

with $\mathcal{K} = 6\ Vol(X_6)$. After introducing the correct Kähler moduli (2.8) for our orientifold compactification X_6, the (inverse) metric G^{ij} for the $h^{(+)}_{(1,1)}(X_6)$ moduli of the $H^{(1,1)}_+(X_6)$ cohomology becomes [28,14]:

$$G^{ij} = -\frac{2}{3}\ \mathcal{K}\ \mathcal{K}^{ij} + 2\ t^i\ t^j \quad , \quad i,j = 1,\ldots,h^{(+)}_{(1,1)}(X_6)\ , \tag{2.43}$$

and the following useful relations hold for the derivatives on the Kähler potential (2.10):

$$K_{T^i} = -2\ \frac{t^i}{\mathcal{K}} \quad , \quad K_{T^i\overline{T}^j} = \frac{G^{ij}}{\mathcal{K}^2}\ . \tag{2.44}$$

Equipped with these relations we may rewrite our no–go criteria (2.36) for a KKLT setup. Since $\mathcal{K} > 0$, we reformulate (2.36) into the condition:

$$\boxed{\forall\ \ i = 1,\ldots,h^{(+)}_{(1,1)}(X_6):\qquad t^i > 0} \tag{2.45}$$

As we have converted the condition (2.36) into (2.45), we shall rewrite the criterion (2.40) in terms of the quantities introduced in (2.41) and (2.42). First, with (2.43) and (2.44) we obtain for the matrix $\mathcal{O}$:

$$\mathcal{O}_{ij} = -K_{T^i\overline{T}^j} + \frac{1}{2}\ K_{T^i}\ K_{\overline{T}^j} = \frac{2}{3}\ \frac{\mathcal{K}^{ij}}{\mathcal{K}}\ .$$

Hence, to obtain the principal minor $\det\mathcal{O}_k$, we essentially have to erase the k–th row and column from the matrix $\mathcal{K}^{ij}$ and compute the resulting determinant. A minor of an inverse matrix A may be computed with the relation [27]:

$$\det A^{-1}(\alpha') = \frac{\det A(\alpha)}{\det A}\ .$$

Here, $A(\alpha)$ denotes the principal submatrix of A, built from the rows and columns α, while $A(\alpha')$ denotes the principal submatrix of A, built from A by deleting the rows and columns α. With this information, we obtain:

$$\det(\mathcal{O}_k) = \left(\frac{2}{3\mathcal{K}}\right)^{n-1} \ \frac{\mathcal{K}_{kk}}{\det(\mathcal{K})} \ .$$

The intersection form $\mathcal{K}_{ij}$ has signature $(1, h^{(+)}_{(1,1)}(X_6)-1)$ [29], *i.e.* $sign(\det\mathcal{K}) = (-1)^{n-1}$. Therefore, the condition $(-1)^n \ \det\mathcal{O}_k \leq 0$ of (2.40) is equivalent to $\mathcal{K}_{kk} \geq 0$ and the criterion (2.40) is translated into:

$$\boxed{\exists \ \ i \in \{\ 1, \ldots, h^{(+)}_{(1,1)}(X_6)\ \}: \quad t^i > 0 \quad \bigwedge \quad \mathcal{K}_{ii} \geq 0} \tag{2.46}$$

Here, $\mathcal{K}_{ii}$ is the i–th diagonal element of the intersection form $\mathcal{K}_{ij}$. To summarize (2.46): if we find one positive Kähler modulus $t^i > 0$ and if in addition we have $\mathcal{K}_{ii} \geq 0$, a stable uplift is not possible with the superpotential (2.23). The latter consists of a sum over contributing divisors $\mathcal{D}_i$ whose Poincaré dual classes ω_i correspond to the Kähler moduli t^i in (2.41). From the relation $\mathcal{K}_i \ t^i = \frac{4}{3} \ Vol(\mathcal{D}_i) \ t^i = \mathcal{K} = 6Vol(X_6) > 0$ and $Vol(\mathcal{D}_i) > 0$, we see, that there *always* exists at least one positive Kähler modulus $t^k > 0$ in the expansion (2.41). Therefore, according to (2.46) we only have to verify whether for this Kähler modulus t^k, we have $\mathcal{K}_{kk} \geq 0$.

The condition (2.45) is the inequality (2.36) written in terms of the Kähler coordinates t^j of J (*cf. Eq.* (2.41)). If all of them are positive, *i.e.* $t^j > 0$, the mass matrix M for the axions is negative definite. For instance this is the case, when the Kähler class J of X_6, defined in (2.41), lies in the Kähler cone, *i.e.* when we have [29]

$$\int_{C_2} J > 0 \quad , \quad \int_{C_4} J \wedge J > 0 \quad , \quad \int_{X_6} J \wedge J \wedge J > 0 \tag{2.47}$$

for all curves C_2 and 4–cycles of the CYM X_6. Then we may define the Kähler coordinates t^j through $t^j = \int_{C_2^j} J > 0$ and meet the criteria (2.45).

To summarize : If (i) the Kähler class J lies in the Kähler cone and (ii) each Kähler modulus t^j corresponds to a 4–cycle of volume $\mathrm{Re}(T^j)$ contributing to (2.23), *no* stable uplift is possible for $h^{+}_{(2,1)}(X_6) = 0$.

To this statement we have to add two refinements for the case that the above two assumptions (i) and (ii) do not hold. Let us first discuss point (i). The Kähler moduli t^j may not be equal to the real coordinates $\widetilde{t^j}$ on the space of Kähler classes of the CY manifold X_6. The latter fulfill the positivity conditions (2.47). Generically, there are linear relations between the supergravity moduli t^j we are working with and the CY moduli $\widetilde{t^j}$. The latter describe deformations of the Kähler form $\mathcal{J} = \sum\limits_{i=1}^{h_{(1,1)}(X_6)} \widetilde{t^i}\ \widetilde{\omega}_i$, which satisfy the positivity conditions (2.47). One may choose Kähler moduli $\widetilde{t^j} := \int_{C_2^j} \mathcal{J} > 0$, while the supergravity moduli t^j may not always be positive. Furthermore, the intersection form $\widetilde{\mathcal{K}}_{ij} = \widetilde{\mathcal{K}}_{ijk}\widetilde{t^k}$ has signature $(1, h_{(1,1)}(X_6) - 1)$.

It may be useful to present an example [3]. Let us consider the Kähler moduli space $\frac{SU(1,1)}{U(1)} \otimes \frac{SU(2,2)}{SU(2)\times SU(2)\times U(1)}$ of the $\mathbf{Z}_4$–orbifold with the Kähler potential:

$$K(t^i, \overline{t}^i) = -\ln\left[\ t^1\ t^2\ t^3 - \frac{1}{2}\ t^3\ (t^4)^2 - \frac{1}{2}\ t^3\ (t^5)^2\ \right]\ . \tag{2.48}$$

We have $\mathcal{K}_{123} = 1, \mathcal{K}_{344} = -1, \mathcal{K}_{355} = -1$ and the intersection form $\mathcal{K}_{ij}$ is:

$$\mathcal{K}_{ij} = \begin{pmatrix} 0 & t^3 & t^2 & 0 & 0 \\ t^3 & 0 & t^1 & 0 & 0 \\ t^2 & t^1 & 0 & -t^4 & -t^5 \\ 0 & 0 & -t^4 & -t^3 & 0 \\ 0 & 0 & -t^5 & 0 & -t^3 \end{pmatrix}\ . \tag{2.49}$$

It may be verified, that the signature of $\mathcal{K}_{ij}$ is indeed $(1,4)$. According (2.8), *i.e.* $\mathrm{Re}T^j = \frac{3}{4}\mathcal{K}_{jkl}\ t^k\ t^l$ the holomorphic coordinates T^i or 4–cycle volumes $\mathrm{Re}T^i$ become:

$$\begin{aligned} \mathrm{Re}(T^1) &= \frac{3}{2}\ t^2\ t^3 \quad , \quad \mathrm{Re}(T^4) = -\frac{3}{2}\ t^3\ t^4\ , \\ \mathrm{Re}(T^2) &= \frac{3}{2}\ t^1\ t^3 \quad , \quad \mathrm{Re}(T^5) = -\frac{3}{2}\ t^3\ t^5\ , \\ \mathrm{Re}(T^3) &= \frac{3}{2}\ \left[\ t^1\ t^2 - \frac{1}{2}\ (t^4)^2 - \frac{1}{2}\ (t^5)^2\ \right]\ . \end{aligned} \tag{2.50}$$

In terms of these moduli, the Kähler potential $-2K(t^i, \overline{t}^i)$ for the orientifold becomes:

$$K(T^i, \overline{T}^i) = -\ln\frac{1}{6\sqrt{6}}\ (T^3 + \overline{T}^3)\left[\ (T^1 + \overline{T}^1)\ (T^2 + \overline{T}^2)\ - \frac{1}{2}\ (T^4 + \overline{T}^4)^2 - \frac{1}{2}\ (T^5 + \overline{T}^5)^2\ \right]\ . \tag{2.51}$$

With (2.49) the equations (2.43) and (2.44) may be verified. The important point here is, that we must have $t^4, t^5 < 0$ for positive 4–cycle volumes $\mathrm{Re}(T^j)$. Hence, in that example the supergravity moduli t^i appearing in (2.41) do not define a Kähler class which lies in the Kähler cone. Nonetheless, though (2.45) is not satisfied, with the Kähler metric (2.51), *i.e.* $K_{KM} = K$, we do not obtain a stable uplift [3].

Let us now come to point (*ii*). The sum of exponentials in the superpotential (2.23) consists of a sum over a basis of $h^{(+)}_{(1,1)}(X_6)$ independent divisor classes $\mathcal{D}_i$ corresponding to the $h^{(+)}_{(1,1)}(X_6)$ 4–cycles the Euclidean $D3$–branes or $D7$–branes are wrapped on. The divisors must have arithmetic genus $\chi(\mathcal{D}_i) = 1$ (*cf.* Section 4) to guarantee that $\gamma_i \neq 0$ in the superpotential (2.23). Let ω_i denote the Poincaré dual class of $\mathcal{D}_i$, which is an element of $H^{(2)}_+(X_6)$. If those classes ω_i are identified with the basis of $(1,1)$–forms in the Kähler class (2.41), the moduli t^j may not be identical to the Kähler deformations $\widetilde{t^i}$ one introduces for the CY orientifold X_6 to begin with. This happens, if the contributing divisor classes $\mathcal{D}_i$ are constructed[4] through linear combinations from the original divisors $\widetilde{\mathcal{D}}_i$, which are Poincaré dual to the classes $\widetilde{\omega}_i$. However, since the basis $\{\omega_i\}$ may be expressed as linear combination of the basis $\{\widetilde{\omega}_i\}$ we obtain linear relations between the two moduli t^j and $\widetilde{t^j}$. When one starts with a Calabi–Yau orientifold X_6 with the Kähler form $\mathcal{J} = \sum\limits_{A=1}^{h_{(1,1)}(X_6)} \widetilde{t^i}\ \widetilde{\omega}_i$ and divisor classes $\widetilde{\mathcal{D}}_i \in H^{(4)}(X_6, \mathbf{Z})$, which are Poincaré dual to the classes $\widetilde{\omega}_i \in H^{(2)}(X_6, \mathbf{Z})$, one first has to find a set of independent divisors $\mathcal{D}_i$ with $\chi(\mathcal{D}_i) = 1$, *i.e.* a linear combination:

$$\mathcal{D}_i = \sum_{j=1}^{h^{(+)}_{(1,1)}(X_6)} n_{ij}\ \widetilde{\mathcal{D}}_j \quad , \quad i = 1, \ldots, h^{(+)}_{(1,1)}(X_6)\ . \tag{2.52}$$

In other words, there exists are an invertible $n \times n$ matrix $N = (n)_{ij}$ such that $\mathcal{D} = N\widetilde{\mathcal{D}}$. The new Kähler coordinates t^i entering the Kähler form (2.41) are given by:

$$\begin{pmatrix} t^1 \\ \vdots \\ t^n \end{pmatrix} = (N^t)^{-1} \begin{pmatrix} \widetilde{t^1} \\ \vdots \\ \widetilde{t^n} \end{pmatrix} \ . \tag{2.53}$$

In terms of the these Kähler coordinates t^j and the basis ω_i, the Kähler form (2.41) becomes a linear combination of contributing divisors $\mathcal{D}_i$ (Poincaré dual to the ω_i). The change of coordinates (2.53) implies the following new intersection form $\mathcal{K}_{ij}$:

$$\mathcal{K}_{ij} = N\widetilde{\mathcal{K}}_{ij}N^t\ . \tag{2.54}$$

Since N is a non–singular matrix, after Sylvester's law of inertia [27], both intersection forms $\mathcal{K}_{ij}$ and $\widetilde{\mathcal{K}}_{ij}$ have the same inertia, *i.e.* they have the signature $(1, h^{(+)}_{(1,1)}(X_6) - 1)$.

---

[4] Note, that this change of basis does not alter the positivity properties of the axionic mass matrix M, introduced in (2.33). In the mass matrix M, the second derivatives are calculated w.r.t. the contributing divisor volumes $\mathrm{Vol}(\mathcal{D}_i)$. The mass matrix w.r.t. the original divisor volumes $\mathrm{Vol}(\widetilde{\mathcal{D}}_i)$ is given by $N^{-1}MN^{\star}$. Since N has rank n the mass matrix $N^{-1}MN^{\star}$ is positive definite if M is positive definite [27].

As an instructive example let us discuss the orientifold of the X_8 CYM, *i.e.* $\mathbf{P}^4[1,1,1,6,9]$ [15]. We may introduce the Kähler form $\mathcal{J} = \widetilde{t}^1\widetilde{\omega}_1 + \widetilde{t}^2\widetilde{\omega}_2$, with the intersection numbers $\widetilde{\mathcal{K}}_{112} = 1, \widetilde{\mathcal{K}}_{122} = 6$ and $\widetilde{\mathcal{K}}_{222} = 36$ and the intersection form $\widetilde{\mathcal{K}}_{ij} = \begin{pmatrix} \widetilde{t}^2 & \widetilde{t}^1 + 6\widetilde{t}^2 \\ \widetilde{t}^1 + 6\widetilde{t}^2 & \widetilde{t}^1 + 36\widetilde{t}^2 \end{pmatrix}$. With this information, the volume of X_6 becomes $Vol(X_6) = \frac{1}{6}[\ 3(\widetilde{t}^1)^2\widetilde{t}^2 + 18\widetilde{t}^1(\widetilde{t}^2)^2 + 36(\widetilde{t}^2)^3\ ]$. According to the formula $V_{\widetilde{\mathcal{D}}_i} = \frac{3}{4}\ \widetilde{\mathcal{K}}_{ijk}\widetilde{t}^j\widetilde{t}^k$ the divisor volumes of the two 4–cycles, whose divisors $\widetilde{\mathcal{D}}_i$ are Poincaré dual to $\widetilde{\omega}_i$ are:

$$\begin{aligned} V_{\widetilde{\mathcal{D}}_1} &= \frac{3}{2}\ \widetilde{t}^2\ (\widetilde{t}^1 + 3\ \widetilde{t}^2)\ , \\ V_{\widetilde{\mathcal{D}}_2} &= \frac{3}{4}\ (\widetilde{t}^1 + 6\ \widetilde{t}^2)^2\ . \end{aligned} \tag{2.55}$$

However, the two divisors $\mathcal{D}_i$ contributing to the superpotential (2.23) are linear combinations (2.52) of the two divisors $\widetilde{\mathcal{D}}_i$ [15], *i.e.*

$$V_{\mathcal{D}_1} = \frac{2}{3}\ (V_{\widetilde{\mathcal{D}}_2} - 6\ V_{\widetilde{\mathcal{D}}_1}) \quad , \quad V_{\mathcal{D}_2} = \frac{2}{3}\ V_{\widetilde{\mathcal{D}}_2}\ , \tag{2.56}$$

with the divisor volumes:

$$\mathrm{Re}(T^1) := V_{\mathcal{D}_1} = \frac{1}{2}\ (\widetilde{t}^1)^2 \quad , \quad \mathrm{Re}(T^2) := V_{\mathcal{D}_2} = \frac{1}{2}\ (\widetilde{t}^1 + 6\ \widetilde{t}^2)^2\ . \tag{2.57}$$

In terms of the latter, the Kähler potential $K_{KM} = -2\ln Vol(X_6)$ for the Kähler moduli T^1, T^2 becomes:

$$K_{KM}(T^j, \overline{T}^j) = \ln 1296 - 2\ln\left[\ (T^2 + \overline{T}^2)^{3/2} - (T^1 + \overline{T}^1)^{3/2}\ \right]\ . \tag{2.58}$$

The divisors $\mathcal{D}_i$, introduced in (2.56), correspond to the 2–forms $\omega_1 = -4\ \widetilde{\omega}_1 + \frac{2}{3}\ \widetilde{\omega}_2$ and $\omega_2 = \frac{2}{3}\ \widetilde{\omega}_2$, respectively. The latter serve as a basis for the Kähler form (2.41), *i.e.* $J = t^1\omega_1 + t^2\omega_2$, with the new[5] Kähler coordinates t^i:

$$t^1 = -\frac{1}{4}\ \widetilde{t}^1 < 0 \quad , \quad t^2 = \frac{1}{4}\ (\widetilde{t}^1 + 6\ \widetilde{t}^2) > 0\ . \tag{2.60}$$

The volume of X_6 may be written $Vol(X_6) = \frac{16}{9}\ [(t^1)^3 + (t^2)^3]$, which gives the intersection numbers $\mathcal{K}_{111} = \mathcal{K}_{222} = \frac{32}{3}$ and the intersection form $\mathcal{K}_{ij} = \frac{32}{3}\ \begin{pmatrix} t^1 & 0 \\ 0 & t^2 \end{pmatrix}$, with signature

---

[5] According to (2.44) these coordinates may be read off from the first derivatives of the Kähler potential $K_{KM} = K$:

$$t^1 = -3V\ K_{T^1} = -\frac{1}{4}\ \widetilde{t}^1 \quad , \quad t^2 = -3V\ K_{T^2} = \frac{1}{4}\ (\widetilde{t}^1 + 6\ \widetilde{t}^2)\ . \tag{2.59}$$

$(1,1)$. Clearly, we have $\mathcal{K}_{ij} = N\widetilde{\mathcal{K}}_{ij}N^t$, with $N = \frac{2}{3}\begin{pmatrix} -6 & 1 \\ 0 & 1 \end{pmatrix}$. Due to (2.60), we see that (2.45) is not fulfilled, *i.e.* this criterion does not give us an answer whether an uplift is possible. However, for the X_8 CYM from above we find:

$$t^2 > 0 \quad , \quad \mathcal{K}_{22} = \frac{32}{3}\, t^2 > 0 \ .$$

According to the second criterion (2.46), no stable uplift would be possible for the X_8 CYM if we assumed the ansatz (2.23) for the superpotential with the two contributing divisors (2.57). From this example we learn, that in practice the inequality (2.45) may not always be satisfied and the condition (2.46) needs to be borrowed.

As we shall see in Subsection 5.3, the condition (2.45) has serious consequences for the resolved orientifolds of orbifolds without complex structure moduli. In fact, for those orbifolds, it is generically not possible to find stable minima in a KKLT setup. This rules out the following orbifolds: $\mathbf{Z}_3, \mathbf{Z}_7, \mathbf{Z}_3 \times \mathbf{Z}_3, \mathbf{Z}_4 \times \mathbf{Z}_4, \mathbf{Z}_6 \times \mathbf{Z}_6, \mathbf{Z}_2 \times \mathbf{Z}_{6'}$, and $\mathbf{Z}_{8-I}$ with $SU(4)^2$ lattice. It would affect even more orbifolds if we would not take into account the twisted complex structure moduli.

2.4. Examples

In the previous subsection we have derived conditions on the Kähler potential $K_{KM} = K$ for the Kähler moduli in type IIB orientifold compactifications without complex structure moduli. These conditions are summarized in *Eqs.* (2.45),(2.46) or alternatively in *Eqs.* (2.36),(2.40). If these criteria are met, a stable uplift in a KKLT setup is not possible, since otherwise the axion mass matrix M would not be positive definite. This way, it may immediately be decided whether a given CY compactification manifold X_6 allows moduli stabilization in the framework of KKLT. In this subsection, we shall apply these conditions to some prominent Kähler potentials K.

For the Kähler potential of the unresolved $\mathbf{Z}_7$ type IIB orientifold [3]

$$K = -\ln(T^1 + \overline{T}^1) - \ln(T^2 + \overline{T}^2) - \ln(T^3 + \overline{T}^3) \tag{2.61}$$

we find:

$$K_{T^1} < 0 \quad , \quad K_{T^2} < 0 \quad , \quad K_{T^3} < 0 \ . \tag{2.62}$$

According to (2.36) the properties (2.62) result in a negative definite axionic mass matrix M.

Furthermore, for the Kähler potential (2.51) of the unresolved $\mathbf{Z}_4$ type IIB orientifold we have:

$$t^1 > 0 \quad , \quad \mathcal{K}_{11} = 0 \ . \tag{2.63}$$

The last relation follows from the intersection form (2.49). Hence due to the condition (2.46), no stable uplift is possible.

Finally, the Kähler potential for the $\mathbf{Z}_3$ type *IIB* orientifold, given in *Eq.* (2.54) of *Ref.* [3] (with the triple intersection numbers (2.56) from there), shares again the property (2.63). Hence, due to the condition (2.46), for the unresolved $\mathbf{Z}_3$–orientifold is no stable uplift possible either.

Note, that the above three examples, which do not allow a stable uplift in the KKLT setup, represent toroidal orbifold/orientifold compactifications, whose stability criteria have been investigated in great detail in [3] (*cf.* also [24]). However, it is intriguing, that we may quickly verify these findings by simply analyzing the conditions (2.36) or (2.40).

A different example is the Kähler potential for the X_8 CYM (2.58), with $(T^2+\overline{T}^2)^{3/2}-(T^1+\overline{T}^1)^{3/2}>0$. Moduli stabilization of the latter has been recently discussed in *Ref.* [30], with the dilaton S and the complex structure moduli being integrated out, *i.e.* $W_0(S,U)=$ *const* in their case. For that case all the equations of Subsection 2.1 apply. In particular, since (*cf.* also *Eq.* (2.59))

$$K_{T^1}=\frac{3}{2}\ \frac{(T^1+\overline{T}^1)^{1/2}}{(T^2+\overline{T}^2)^{3/2}-(T^1+\overline{T}^1)^{3/2}}>0\ ,\ K_{T^2}=-\frac{3}{2}\ \frac{(T^2+\overline{T}^2)^{1/2}}{(T^2+\overline{T}^2)^{3/2}-(T^1+\overline{T}^1)^{3/2}}<0 \tag{2.64}$$

from (2.21) it follows, $\phi_{12}=a_1t_2^1-a_2t_2^2=\pi$, which indeed has been verified in [30]. On the other hand, if we used the Kähler potential (2.58) together with the superpotential (2.23) we would immediately see, that in this case no stable uplift would be possible. This may be easily seen from condition (2.40). We find:

$$K_{T^2}<0\quad ,\quad -K_{T^1\overline{T}^1}+\frac{1}{2}\ K_{T^1}^2=-\frac{3}{8}\ \frac{(T^1+\overline{T}^1)^{3/2}+2\ (T^2+\overline{T}^2)^{3/2}}{(T^1+\overline{T}^1)^{1/2}\ [(T^2+\overline{T}^2)^{3/2}-(T^1+\overline{T}^1)^{3/2}]^2}<0\ . \tag{2.65}$$

3. Stabilization of Kähler moduli associated to the cohomology $H_-^{(1,1)}(X_6)$

Because the superpotential W does not depend on the $h_{(1,1)}^{(-)}(X_6)$ moduli, *i.e.* $\partial_{G^a}W=0$, the whole KKLT stabilization procedure only works for the $h_{(1,1)}^{(+)}(X_6)$ Kähler moduli T^k, while we have to find a different mechanism to stabilize the 2 $h_{(1,1)}^{(-)}(X_6)$ real fields b^a, c^a, which are combined into the $h_{(1,1)}^{(-)}(X_6)$ N=1 chiral fields (2.5). Since the resolved $\mathbf{Z}_N$ and $\mathbf{Z}_N\times\mathbf{Z}_M$ orbifolds, whose vacuum structure will be the main part of Section 4, generically have those moduli G^a (*cf.* Table 4), fixing them becomes an important question for these vacua.

A first look at the F–term potential

$$V_F = K_{I\overline{J}}\ F^I \overline{F}^{\overline{J}} - 3\ e^{\kappa_4^2 K}\ \kappa_4^2\ |W|^2\ , \tag{3.1}$$

with $I, \overline{J}$ running over all fields S, T^j, G^a, U^λ and the vacuum conditions

$$F^{T^j} = 0 \quad , \quad F^{G^a} = 0 \quad , \quad F^S = 0 \quad , \quad F^{U^\lambda} = 0$$

in the full moduli space yields ($\partial_{G^a} W = 0$):

$$\begin{aligned} \partial_S W + W\ \partial_S K &= 0\ , \\ \partial_{T^j} W + W\ \partial_{T^j} K &= 0 \quad , \quad j = 1, \ldots, h^{(+)}_{(1,1)}(X_6)\ , \\ W\ K_{G^a} &= 0 \quad , \quad a = 1, \ldots, h^{(-)}_{(1,1)}(X_6)\ , \\ \partial_{U^\lambda} W + W\ \partial_{U^\lambda} K &= 0 \quad , \quad \lambda = 1, \ldots, h^{(-)}_{(2,1)}(X_6)\ . \end{aligned} \tag{3.2}$$

As demonstrated in *Refs.* [14], the Kähler derivatives $\partial_{T^j} K, \partial_S K$ and $\partial_{U^\lambda} K$ become non–trivial functions depending on all moduli T^j, G^a, S. Since at the AdS minimum $W \neq 0$, it follows that

$$K_{G^a} = 0\ . \tag{3.3}$$

According to [14], there is the relation

$$K_{G^a} = -\frac{3}{2}\ K_{T^j}\ \mathcal{K}_{jac} b^c = 3\ \mathcal{K}^{-1}\ \mathcal{K}_{jac} t^j\ b^c = 3\ \mathcal{K}^{-1}\ \mathcal{K}_{ac}\ b^c\ .$$

Therefore, the solution to (3.2) yields:

$$\mathcal{K}_{ac}\ b^c = 0 \quad , \quad a = 1, \ldots, h^{(-)}_{(1,1)}(X_6)\ . \tag{3.4}$$

Thanks to this condition, to be satisfied at the AdS minimum, all the single derivatives $\partial_S K$ and $\partial_{U^\lambda} K$ become the usual ones, and the F–term conditions for the dilaton and complex structure moduli become the usual ones. Hence we may proceed with the stabilization procedure as outlined before. The condition (3.4) may be solved with $b^a = 0$, *i.e.* half of the degrees of freedom of the fields $G^a = g_1^a + i g_2^a$ may be fixed. However, after calculating the scalar potential and the masses for the moduli fields it may be seen, that the masses for the g_1^a fields become negative

$$m_{g_1^a g_1^a} = -e^K\ |W|^2\ \frac{\partial^2 K}{\partial (g_1^a)^2}\ , \tag{3.5}$$

with $\frac{\partial^2 K}{\partial (g_1^a)^2} = 4 K_{G^a \overline{G}^a}$, while the masses for the fields g_2^a are zero:

$$m_{g_2^a g_2^a} = 0\ . \tag{3.6}$$

The latter observation is clear, since neither the Kähler potential nor the superpotential do depend on them. Therefore, the scalar potential does not depend on them either. Hence, two problems have occurred: First, the fields g_2^a cannot be stabilized through the F–term equations (3.2). Second, though the fields g_1^a are stabilized, their masses are negative. In fact, it has been recently shown in *Ref.* [31], that the second problem follows from the first problem. However, this situation is improved in the presence of additional D–terms, which we shall discuss in the following.

In the case of $h^{(-)}_{(1,1)}(X_6) \neq 0$ in the geometry Y_6 of the covering space of the orientifold theory, some divisors E (or divisor orbits under the orbifold group) are not invariant under the orientifold action σ, but are mapped to other divisors $\widetilde{E}$ (or orbits under the orbifold group), *i.e.* :

$$\sigma E = \widetilde{E} \ . \tag{3.7}$$

In that case, invariant combinations have to be constructed[6]. Let

$$E_i := \frac{1}{2}\ (E + \widetilde{E}) \quad , \quad E_a := \frac{1}{2}\ (E - \widetilde{E}) \tag{3.8}$$

such that:

$$\sigma E_i = E_i \quad , \quad \sigma E_a = -E_a \ . \tag{3.9}$$

Then we have $\omega_i \in H^{(1,1)}_+(X_6)$ and $\omega_a \in H^{(1,1)}_-(X_6)$ for their corresponding Hodge dual 2–forms. This way, the $h^{(-)}_{(1,1)}(X_6)$ odd forms ω_a are paired with $h^{(-)}_{(1,1)}(X_6)$ even forms ω_i. After the discussion in Subsection 2.1, a non–vanishing $h^{(-)}_{(1,1)}(X_6) \neq 0$ implies a non–trivial bulk $NSNS$ 2–form B_2

$$B_2 = \sum_{\widetilde{a}=1}^{h^{(-)}_{(1,1)}(X_6)} b^{\widetilde{a}}\ \omega_{\widetilde{a}}\ , \tag{3.10}$$

which (more precisely its pullback on C_i) enters the calibration conditions of a holomorphic 4–cycle C_i on which a $D3$ or $D7$–brane is wrapped upon [32]. If one supersymmetry should be preserved, those conditions are solved by demanding [33]

$$B_2 \wedge J = 0 \ . \tag{3.11}$$

Let us assume the 4–cycle C_i to correspond to the even divisor E_i and the 4–cycle C_a to correspond to the odd divisor E_a. Then, integrating (3.11) over the odd cycle C_a yields:

$$\int_{C_a} B_2 \wedge J = t^j\ b^{\widetilde{a}} \int_{X_6} \omega_{\widetilde{a}} \wedge \omega_j \wedge \omega_a = \mathcal{K}_{ja\widetilde{a}}\ t^j\ b^{\widetilde{a}} = \mathcal{K}_{\widetilde{a}a}\ b^{\widetilde{a}} = 0 \ . \tag{3.12}$$

---

[6] As an illustrative example see the $\mathbf{Z}_{6-II}$ orbifold in Subsection 5.4.

Hence, we conclude:

$$\mathcal{K}_{\widetilde{a}a}\ b^{\widetilde{a}} = 0\ . \tag{3.13}$$

The calibration condition (3.11) has to be imposed on all even 4–cycles C_i. Therefore the condition (3.11) gives rise to a system of $h^{(-)}_{(1,1)}(X_6)$ linear equations for the $h^{(-)}_{(1,1)}(X_6)$ variables b^a. The intersection form $\mathcal{K}_{ab}$ depends also on the $h^{(+)}_{(1,1)}(X_6)$ Kähler moduli T^j. However, generically this matrix has non–vanishing determinant and the only solution to (3.13) is the trivial solution[7]:

$$\mathrm{Re}(G^a) = 0 \quad \Longrightarrow \quad b^a = 0 \quad , \quad a = 1, \ldots h^{(-)}_{(1,1)}(X_6)\ . \tag{3.14}$$

Interestingly, the requirement (3.12) is similar to the condition (3.4) following from the F–terms.

For a $D3$–brane wrapped on the 4–cycle C_i, the requirement (3.11) allows to rewrite the real part of the bosonic instanton action

$$e^{-\phi_{10}} \int_{C_i} d^4\xi\ \sqrt{\det(G+B)} = \frac{1}{2} \int_{C_i} J \wedge J - e^{-\phi_{10}}\ B \wedge B = \frac{2}{3}\ \mathrm{Re}(T^i)\ , \tag{3.15}$$

with T^i given in (2.8) (and the internal CY metric G).

On the other hand, for a stack of $D7$–branes wrapped on the 4–cycle C_i, the requirement (3.13) may be also anticipated from the D–term potential [33]

$$V_D^i \sim \frac{1}{\mathrm{Re}(T^i)}\ (\ \mathcal{K}_{a\widetilde{a}}\ b^{\widetilde{a}}\ )^2\ , \tag{3.16}$$

associated to a $U(1)$ gauge group[8] whose gauge coupling is given by $g^{-2} = \mathrm{Re}(T^i)$ (the $D7$–brane associated to this $U(1)$ is assumed to wrap the 4–cycle C_i). For a stack of $D7$–branes with $U(N)$ gauge group, the remaining $SU(N)$ gauge group may furnish gaugino condensation. The potential (3.16) is minimized if condition (3.12) holds.

Let us now come to the axionic moduli c^a in (2.5). The case $h^{(-)}_{(1,1)}(X_6) \neq 0$ implies the non–trivial bulk RR 2–form C_2:

$$C_2 = \sum_{a=1}^{h^{(-)}_{(1,1)}(X_6)} c^a\ \omega_a\ . \tag{3.17}$$

---

[7] As an illustrative example see the $\mathbf{Z}_{6-II}$ orbifold in Subsection 5.4.

[8] In the case of additional matter charged under this $U(1)$, the potential (3.16) receives an additional term proportional to the matter fields C_i. Such matter fields originate from open strings stretched between different $D7$ or/and $D3$–branes. For simplicity we shall assume that these matter fields are minimized at $C_i = 0$.

If the stack of $D7$–branes wrapping the 4–cycle C_i has a gauge group with an $U(1)$–factor, the fields c^a and ρ^i are gauged under this $U(1)$ gauge field A_μ as a result of the $D7$ world–volume couplings $C_6 \wedge F_2$ and $C_4 \wedge B_2 \wedge F_2$, respectively, producing Green–Schwarz terms. As a consequence of the definition (2.5) the field G^a is charged under this $U(1)$, while the Kähler modulus T^i remains neutral due to a non–trivial cancellation of the gauge transformations of the axionic fields c^a and ρ^i entering the definition of T^i (2.8), *i.e.*

$$D_\mu G^a = \partial_\mu G^a - 4\ i\ \kappa_4^2\ \mu_7\ (2\pi\alpha')\ A_\mu \quad , \quad D_\mu T^j = \partial_\mu T^j\ , \tag{3.18}$$

cf. Ref. [33] for more details. Hence, the D–term potential (3.16) stabilizes the fields b^a, while the covariant derivative $D_\mu G^a$ gives rise to a massive $U(1)$ vector A_μ thus fixing the imaginary part of G^a, *i.e.* the field c^a [34]. The latter is absorbed into the mass of the gauge field. Since only $\mathrm{Im}(T^j)$, but not $\mathrm{Im}(G^a)$ couples to any $F \wedge F$ piece, the mass term may even occur without an anomaly as a Stückelberg mass term [35,36]. To summarize so far, the calibration condition (3.11) and the gauging (3.18) of the modulus G^a allow to stabilize the Kähler moduli associated to the cohomology $H_-^{(1,1)}(X_6)$.

The mechanism to stabilize the axionic fields c^a relies on the existence of a $U(1)$. If the stack of $D7$–branes sits on an orientifold plane we encounter an SO– or Sp gauge group and the previous mechanism has to be modified. In that case, we turn on additional world–volume 2–form fluxes on that stack of $D7$–branes to break off one $U(1)$ factor. In the orientifold on a stack of $D7$–branes, there are two kinds of world–volume 2–form fluxes ${}^Y\!f$ and $\widetilde{f}$. The flux ${}^Y\!f$ is inherited from the ambient CY space Y, while the flux $\widetilde{f}$ is a harmonic 2–form of the 4–cycle C_i. We refer the reader to *Ref.* [33] for a description of the orthogonal splitting of a general 2–form flux f into $f = {}^Y\!f + \widetilde{f}$, with ${}^Y\!f \in im(\iota^\star)$, $\widetilde{f} \in coker(\iota^\star)$, with the map $\iota^\star : H_-^{(2)}(Y_6) \to H_-^{(2)}(C_i)$. In the following, we do not want to consider the class of fluxes $\widetilde{f}$, since they would gauge the Kähler moduli T^j, *i.e.* $\widetilde{f} = 0$. On the other hand, turning on 2–form fluxes ${}^Y\!f_2$ from the ambient space Y_6,

$${}^Y\!f_2 = \sum_{a=1}^{h_{(1,1)}^{(-)}(Y_6)} f^a\ \iota^\star \omega_a, \tag{3.19}$$

does not gauge the Kähler moduli T^j [33]. These fluxes only modify the previous equations with the effect, that the 2–form B_2 is replaced by the combination $B_2 - 2\pi\alpha'\ {}^Y\!f_2$, *i.e.* we supplement *Eqs.* (2.5),(3.11), (3.13), (3.15) and (3.16) with the following substitution [33]:

$$B_2 \longrightarrow B_2 - 2\pi\alpha'\ {}^Y\!f_2 \quad , \quad b^a \longrightarrow b^a - 2\pi\alpha'\ {}^Y\!f^a\ . \tag{3.20}$$

Hence the calibration condition (3.11) is changed into

$$\mathcal{K}_{\widetilde{a}a}\ \left(\ b^{\widetilde{a}} - 2\pi\alpha'\ {}^Y\!f^{\widetilde{a}}\ \right) = 0\ , \tag{3.21}$$

instead of (3.12). Therefore, the solution (3.14) becomes:

$$b^a = 2\pi\alpha' \ {}^Y\!f^a \quad , \quad a = 1, \ldots h^{(-)}_{(1,1)}(X_6) \ . \tag{3.22}$$

Since the gaugings (3.18) now hold for the redefined modulus

$$G^a = i \ c^a - S \ (b^a - 2\pi\alpha' \ {}^Y\!f^a),$$

the previous discussion about the stabilization of the axion c^a by making one $U(1)$ gauge field A_μ massive may be directly applied.

To conclude, the calibration condition (3.11) together with the gauging (3.18) of the modulus G^a allow to stabilize this modulus by making one $U(1)$ gauge field A_μ massive. If the latter does not exist, it may be generated by turning on 2–form fluxes ${}^Y\!f$ from the ambient space Y. It is crucial, that under the background fluxes b^a and ${}^Y\!f^a$ the Kähler modulus T^i remains uncharged, *i.e.* (3.18) holds.

In N=2 supergravity, background fluxes may be associated to some gaugings of continuous PQ–symmetries acting on axions of the hypermultiplet scalars. In the type IIB case, the 3–form fluxes G_3 correspond to gaugings of scalars of the universal hypermultiplet, while the fields c^a and ρ^i are gauged by turning on non–trivial b^a–fields and the world volume 2–form fluxes ${}^Y\!f^a, \widetilde{f}^a$ on the D–brane world volume. Fluxes may protect those isometries, which are gauged in supergravity, from quantum corrections. A kind of complementarity arises between turning on fluxes, *i.e.* gauging some continuous PQ symmetries of hypermultiplet scalars and instanton corrections, which break continuous PQ symmetries. The compatibility of both effects has been recently studied in [37]. After turning on world volume 2–form fluxes, the axions of the Kähler moduli T^i and G^a are the potential candidates for being gauged. Remember, that the Kähler moduli T^i correspond to contributing divisors in the non–perturbative superpotential (2.23) accounting for $D3$–instanton effects and gaugino condensation. Hence, their continuous PQ symmetries are broken by instanton effects rather than by fluxes. However, in the case that only the flux ${}^Y\!f$ from the ambient space Y is turned on, while the 2–form flux $\widetilde{f}$ is absent, the Kähler modulus T^i indeed remains neutral [33]. It is that case we have focused on in the above discussion. Hence the instanton effects in the KKLT superpotential (2.23) are compatible with turning on only 2–form fluxes ${}^Y\!f^a$ from the ambient space Y. We have a complementary situation: one class of Kähler moduli T^i is stabilized through instanton effects in the superpotential, while the other class of Kähler moduli G^a is fixed through gauging isometries (3.18) and turning on the background fluxes $b^a, {}^Y\!f^a$.

4. Instanton effects

In this section, we discuss the question for which cases a non–perturbative superpotential from brane instantons is produced. As shown in [9], if a divisor, wrapped by a $M5$-brane in the dual M-theory picture (dual to type IIB which one is considering), has holomorphic Euler characteristic

$$\chi = h_{(0,0)} - h_{(0,1)} + h_{(0,2)} - h_{(0,3)} = 1,$$

the necessary two fermionic zero modes for the instanton contribution will be present. To discuss our models we use the results of [38–42], where it was shown how the zero modes counting is changed in the presence of background fluxes. The advantage of the counting procedure in [41] is that it is not necessary to do an F-theory lift, the calculations can be done directly in the type IIB picture. Additionally, it was shown in [43] that only the $(2,1)$–component of the G_3-flux may lift zero modes.

To calculate the number of the zero modes, we have to realize what are the possible 4-cycles wrapped by the $D3$-branes in the compact space. The Hodge numbers $h_{(0,0)}, h_{(0,1)}, h_{(0,2)}$ of the 4-cycle give the number of the zero modes with positive (N_+) and negative (N_-) chirality with respect to the normal bundle of the $D3$–brane. If one takes into account background fluxes, orientifold action and fixing of the κ-symmetry, some of the zero modes could be lifted and the index

$$\chi_{D3} = \frac{1}{2}\left(N_+ - N_-\right)$$

will change. χ_{D3} is not anymore of purely geometrical nature. In the case of type IIB, Bergshoeff et al. [41] showed that only $h_{(0,1)}$ and $h_{(0,2)}$ of N_+ can be lifted by fluxes. Thus, if the topology of the divisor has vanishing $h_{(0,1)}$, $h_{(0,2)}$, we can neglect the effect of the fluxes altogether and concentrate only on the action of the O–planes on the zero mode counting.

The correspondence between zero modes of the Dirac operator on the worldvolume of the 4-cycles and Hodge numbers $h_{(0,0)}, h_{(0,1)}, h_{(0,2)}$ of these cycles becomes apparent by mapping the spinors to $(0,p)$-differential forms.[9] Then fermionic zero modes of the Dirac operator correspond to the harmonic forms by above mapping. Locally we can write the world volume spinors on the $D3$–brane as

$$\begin{aligned} \epsilon_+ &= \phi|\Omega> + \phi_{\bar a}\gamma^{\bar a}|\Omega> + \phi_{\overline{ab}}\gamma^{\overline{ab}}|\Omega> \ , \\ \epsilon_- &= \phi_{\bar z}\gamma^{\bar z}|\Omega> + \phi_{\overline{az}}\gamma^{\overline{az}}|\Omega> + \phi_{\overline{abz}}\gamma^{\overline{abz}}|\Omega> \ , \end{aligned} \qquad (4.1)$$

---

[9] $\phi_{a_1\ldots a_N}\gamma^{a_1\ldots a_N}|\Omega> \longleftrightarrow \phi_{a_1\ldots a_N} dz^{a_1}\ldots dz^{a_N}$

where ϵ_+ and ϵ_- are states with positive and negative chirality with respect to the normal bundle $SO(2)$ of the $D3$–brane inside the compact space. a, b are the $D3$–brane worldvolume directions, z is the normal direction to the worldvolume.

Note that ϵ_+ and ϵ_- transform under $SO(4) \times SO(2) \times SO(1,3)$ and the modes ϕ have an additional spinor index which transforms in the $\mathbf{2} \oplus \bar{\mathbf{2}}$ under $SO(1,3)$. Thus, the number of the zero modes given by the Hodge numbers of the 4-cycle has to be doubled.

All modes of ϵ_- have legs in the normal direction to the $D3$–brane. By use of Serre's generalization of the Poincaré duality, these modes can be mapped to those taking values in the worldvolume of $D3$–brane. This duality maps $(0,p)$-forms with values in the bundle $\Omega^{0,p}(X)$ of the 4–cycle X to $(0, 2-p)$–forms with values in $\Omega^{(0,2-p)} \otimes K$, where K is the canonical bundle of the 4–cycle. In the case of the wrapped $D3$–brane, the canonical bundle is equal to the normal bundle, so this duality is realized by multiplying by the covariantly constant 3–form Ω_{abc} and building the Hodge[10] dual.

$$\begin{aligned} g^{z\bar{z}}\Omega_{\overline{abz}}\phi_z &= \widetilde{\phi}_{\overline{ab}} \ , \\ g^{z\bar{z}}g^{a\bar{a}}\Omega_{\overline{abz}}\phi_{az} &= \widetilde{\phi}_{\overline{a}} \ , \\ g^{z\bar{z}}g^{a\bar{a}}g^{b\bar{b}}\Omega_{\overline{abz}}\phi_{abz} &= \widetilde{\phi} \ . \end{aligned} \tag{4.3}$$

This means that the numbers of the modes with positive and negative chirality match. If all zero modes are present, the corresponding index

$$\chi_{D3} = \frac{1}{2}\ (N_+ - N_-) = \big(\ h_{(0,0)} + h_{(0,1)} + h_{(0,2)}\ \big) - \big(\ h_{(0,0)} + h_{(0,1)} + h_{(0,2)}\ \big) \tag{4.4}$$

will be 0.

4.1. Calculation of χ_{D3} for divisors with $h_{(0,0)} = 1$, $h_{(0,1)} = h_{(0,2)} = 0$

As we shall see in the next section, many of the divisors arising in resolved toroidal orbifold models have the Hodge numbers $h_{(0,0)} = 1$, $h_{(0,1)} = h_{(0,2)} = 0$. We will therefore start by calculating the number of zero modes for this especially simple case. We choose $a, \bar{a}, b, \bar{b}$ as holomorphic and antiholomorphic coordinates on the $D3$–brane, which take the values $1, \bar{1}, 2, \bar{2}$. z and $\bar{z}$ should correspond to the transverse directions with values $3, \bar{3}$.

---

[10]

$$\begin{aligned} &*\Big(\phi_{\overline{a}_1\dots\overline{a}_{N-p}}\Omega_{a_1\dots a_N}\,dz^{\overline{a}_1}\dots dz^{\overline{a}_{N-p}}dz^{a_1}\dots dz^{a_N}\Big) \\ &\qquad = \epsilon_{\overline{a}_1\dots\overline{a}_N}\epsilon_{a_1\dots a_N}\phi^{\overline{a}_1\dots\overline{a}_{N-p}}\Omega^{a_1\dots a_N}\ dz^{\overline{N-p+1}}\dots dz^{\overline{N}} \end{aligned} \tag{4.2}$$

Note, that in our convention the form is complex conjugated by applying the Hodge star.

The fermionic states on the $D3$–brane corresponding to $h_{(0,0)}$ are

$$\epsilon_+ = \phi|\Omega> \ , \qquad \epsilon_- = \phi_{\overline{abz}}\gamma^{\overline{abz}}|\Omega> \ . \tag{4.5}$$

On the brane, some of the modes are pure gauge due to the κ-symmetry. These are the modes which are annihilated by the κ-symmetry projector $(1-\Gamma_{D3})\theta = 0$, where $\Gamma_{D3} = \sigma_2 \otimes \gamma_5$ with γ_5 four ten dimensional γ-matrices pulled back on the brane. θ corresponds to two 32–component spinors written in the double spinor formalism [44]. Additionally, some of the modes can be projected out by the orientifold action. We have to choose κ-symmetry fixing in such a way that it commutes with the orientifold action [41].

There are three different cases to distinguish for the position of the $O7$–plane: it can be on top of the D3–brane, can intersect it along one direction, or can be parallel to it. We assume that the $O7$–plane fills the non-compact directions.

- Case 1: an O-plane lies on top of a D3–brane

It is convenient to do the calculations in the local coordinate patch. The κ-symmetry fixing condition and the projection through the orientifold action are given by

$$\begin{aligned}(1-\sigma_2\gamma^{1\bar{1}2\bar{2}})\theta = 0 \ , \\ (1-\sigma_2\gamma^{3\bar{3}})\theta = 0 \ .\end{aligned} \tag{4.6}$$

Both conditions written together yield

$$(1-\gamma^{1\bar{1}2\bar{2}3\bar{3}})\theta = 0 \ . \tag{4.7}$$

Inserting $\theta = \epsilon_+ + \epsilon_-$ shows that ϕ survives this projection and $\phi_{\overline{abz}}$ not. The index is $\chi_{D3} = h_{(0,0)} = 1$.

- Case 2: Intersection with an O–plane along one complex dimension

The O–plane intersects the D–brane along complex direction 1. Then, κ-symmetry fixing condition and the projection through the orientifold action are given by

$$\begin{aligned}(1-\sigma_2\gamma^{1\bar{1}2\bar{2}})\theta = 0 \ , \\ (1-\sigma_2\gamma^{2\bar{2}})\theta = 0 \ .\end{aligned} \tag{4.8}$$

Both conditions written together give

$$(1-\gamma^{1\bar{1}})\theta = 0 \ . \tag{4.9}$$

ϕ survives this projection, $\phi_{\overline{abz}}$ not. From this it follows $\chi_{D3} = h_{(0,0)} = 1$.

- Case 3: No intersection with an O–plane

It can be the case, when the O–plane is parallel to the $D3$–brane. The orientifold action maps fermions of the brane to the fermions in the mirror brane, so no modes are projected out. There is only the κ-symmetry fixing condition, by which no modes are cut. The modes $\phi|\Omega>$, $\phi_{\overline{abc}}\gamma^{\overline{abc}}|\Omega>$ corresponding to $h_{(0,0)}$ are present and the index is $\chi_{D3} = h_{(0,0)} - h_{(0,0)} = 0$.

By investigating all configurations of the $O7$–plane we obtain a general statement:

Divisors with Hodge numbers $h_{(0,0)} = 1$, $h_{(0,1)} = h_{(0,2)} = 0$ *will have the index* $\chi_{D3} = 1$ *if an* $O7$*–plane lies on top of them or if it intersects the divisor along one complex dimension. Otherwise,* $\chi_{D3} = 0$.

4.2. General case: $h_{(0,1)}, h_{(0,2)} \neq 0$

As discussed in the last subsection, locally, there are always only three different configurations of the $O7$–plane relative to the divisor in question. It can be on top of it, intersect it in one complex direction, or be parallel to it. a, b are again the coordinates on the $D3$–brane. The projector equations from the fixing of the κ-symmetry and the orientifold action will be as in the previous subsection. The only difference is that the modes $\phi_{\overline{a}}|\Omega>$ and $\phi_{\overline{ab}}\gamma^{\overline{ab}}|\Omega>$ are now present. They can be lifted by fluxes. When turning on fluxes, we assume that they will be of the the most unfavorable form for the presence of zero modes. This would correspond to a general form for the fluxes. We summarize the results of the action of the projector equations in all three cases in the following table:

	$O-$ plane on top of $D3$		$O-$ plane intersects $D3$		$O-$ plane does not intersect $D3$	
chirality	$+$	$-$	$+$	$-$	$+$	$-$
$h_{(0,0)}$	ϕ		ϕ		ϕ	$\phi_{\overline{abz}}$
$h_{(0,1)}$		$\phi_{\overline{az}}$	$[\phi_{\overline{a}}]$	$\phi_{\overline{az}}$	$[\phi_{\overline{a}}]$	$\phi_{\overline{az}}$
$h_{(0,2)}$	$[\phi_{\overline{ab}}]$			$\phi_{\overline{z}}$	$[\phi_{\overline{ab}}]$	$\phi_{\overline{z}}$
# of zero modes	$2 - 2\ h^{(-)}_{(0,1)} + 2\ [h^{(+)}_{(0,2)}]$		$2 - 2\ h^{(-)}_{(0,1)}$ $-2\ h^{(-)}_{(0,2)} + 2\ [h^{(+)}_{(0,1)}]$		$2\ [h^{(+)}_{(0,1)}] + 2\ [h^{(-)}_{(0,2)}]$ $-2\ h^{(-)}_{(0,1)} - 2\ h^{(+)}_{(0,2)}$	

Table 2: *Zero modes after fixing* κ*-symmetry and orientifold projection*

In the horizontal line we give the zero modes associated to the Hodge numbers $h_{(0,0)}, h_{(0,1)}, h_{(0,2)}$. '+' and '-' denote the chirality with respect to the normal bundle of the $D3$–brane. In brackets we put the modes which are in general lifted in the presence of fluxes, and in the last line we give the number of zero modes which are left.

Let us discuss this result first before turning on flux. The first column represents the case where the influence of the orientifold projection is fully felt by the divisor in question. As in the M–theory case discussed by Witten, only one chirality survives for each Hodge number and the index reduces again to the holomorphic Euler characteristic $h_{(0,0)} - h_{(0,1)} + h_{(0,2)}$. The third column corresponds to the case where the influence of the orientifold is not felt at all by the wrapped divisor. Both chiralities survive and cancel each other out. This agrees with the observation that for a compactification on a CY manifold (without orientifold projection), no non–perturbative superpotential is generated. The second column represents an intermediate case. It is obvious that the knowledge of the Hodge numbers is of prime importance to be able to decide whether a divisor contributes to the non–perturbative superpotential. We can see that without turning on flux, we can get a contribution in the first column for $h_{(0,1)} = h_{(0,2)}$. With flux, a contribution is only possible for $h_{(0,1)} = 0$. We get a contribution from the second column for $h_{(0,2)} = 0$ if no flux is turned on, with flux only for $h_{(0,1)} = h_{(0,2)} = 0$. With or without flux, column three never gives a contribution since the number of zero modes is always less than or equal to zero.

In the present work we do not discuss the counting of zero modes for the case with non–vanishing 2–form flux f on the D–brane world–volume, as it has been discussed in Section 3. Recently, work towards this direction has been accomplished in *Ref.* [45] for the case of heterotic M-theory. The authors have found that world–volume flux does not change the zero mode counting for the case of some particular background fluxes. Those fluxes were chosen such that they do not lift any zero modes. For the case of IIB, Bandos and Sorokin derived the Dirac equation for the $D3$–brane in the presence of worldvolume flux [46]. Its implication for the zero mode counting has to be analyzed [47]. Compared to the case without 2–form flux, there are more complicated conditions on the gauge fixing of the κ-symmetry and an additional field equation for the 2–form flux, which depend on the topology of the Calabi–Yau manifold.

We finish this section by the remark that the formalism described above requires the wrapped 4–cycle to be Kähler. In our models it is the case because we have to deal with divisors which are hyperplanes in the Calabi–Yau manifolds and hyperplanes of a Kähler manifold are Kähler. This can be also directly observed from the topology of the divisors, which are either products or fibrations of tori and $\mathbf{P}^1$s.

5. Fixing all moduli in resolved type IIB $\mathbf{Z}_N-$ and $\mathbf{Z}_N \times \mathbf{Z}_M$–orientifolds

5.1. Type IIB orientifolds of $\mathbf{Z}_N-$ *and* $\mathbf{Z}_N \times \mathbf{Z}_M$*–orbifolds*

In the following we shall investigate moduli stabilization for a class of type IIB orientifold compactifications X_6. We shall discuss orientifolds X_6 of the resolved toroidal orbifolds Y_6

$$Y_6 = T^6/\Gamma \quad , \quad \Gamma = \mathbf{Z}_N \quad , \quad \mathbf{Z}_N \times \mathbf{Z}_M \ , \tag{5.1}$$

with orbifold group Γ.

$\mathbf{Z}_N$	lattice T^6	$h^{\text{untw.}}_{(1,1)}$	$h^{\text{untw.}}_{(2,1)}$	$h^{\text{twist.}}_{(1,1)}$	$h^{\text{twist.}}_{(2,1)}$
$\mathbf{Z}_3$	$SU(3)^3$	9	0	27	0
$\mathbf{Z}_4$	$SU(4)^2$	5	1	20	0
$\mathbf{Z}_4$	$SU(2) \times SU(4) \times SO(5)$	5	1	22	2
$\mathbf{Z}_4$	$SU(2)^2 \times SO(5)^2$	5	1	26	6
$\mathbf{Z}_{6-I}$	$(G_2 \times SU(3)^2)^\flat$	5	0	20	1
$\mathbf{Z}_{6-I}$	$SU(3) \times G_2^2$	5	0	24	5
$\mathbf{Z}_{6-II}$	$SU(2) \times SU(6)$	3	1	22	0
$\mathbf{Z}_{6-II}$	$SU(3) \times SO(8)$	3	1	26	4
$\mathbf{Z}_{6-II}$	$(SU(2)^2 \times SU(3) \times SU(3))^\sharp$	3	1	28	6
$\mathbf{Z}_{6-II}$	$SU(2)^2 \times SU(3) \times G_2$	3	1	32	10
$\mathbf{Z}_7$	$SU(7)$	3	0	21	0
$\mathbf{Z}_{8-I}$	$(SU(4) \times SU(4))^*$	3	0	21	0
$\mathbf{Z}_{8-I}$	$SO(5) \times SO(9)$	3	0	24	3
$\mathbf{Z}_{8-II}$	$SU(2) \times SO(10)$	3	1	24	2
$\mathbf{Z}_{8-II}$	$SO(4) \times SO(9)$	3	1	28	6
$\mathbf{Z}_{12-I}$	E_6	3	0	22	1
$\mathbf{Z}_{12-I}$	$SU(3) \times F_4$	3	0	26	5
$\mathbf{Z}_{12-II}$	$SO(4) \times F_4$	3	1	28	6
$\mathbf{Z}_2 \times \mathbf{Z}_2$	$SU(2)^6$	3	3	48	0
$\mathbf{Z}_2 \times \mathbf{Z}_4$	$SU(2)^2 \times SO(5)^2$	3	1	58	0
$\mathbf{Z}_2 \times \mathbf{Z}_6$	$SU(2)^2 \times SU(3) \times G_2$	3	1	48	2
$\mathbf{Z}_2 \times \mathbf{Z}_{6'}$	$SU(3) \times G_2^2$	3	0	33	0
$\mathbf{Z}_3 \times \mathbf{Z}_3$	$SU(3)^3$	3	0	81	0
$\mathbf{Z}_3 \times \mathbf{Z}_6$	$SU(3) \times G_2^2$	3	0	70	1
$\mathbf{Z}_4 \times \mathbf{Z}_4$	$SO(5)^3$	3	0	87	0
$\mathbf{Z}_6 \times \mathbf{Z}_6$	G_2^3	3	0	81	0

Table 3: *Orbifold groups, lattices and Hodge numbers for* $\mathbf{Z}_N$ *and* $\mathbf{Z}_N \times \mathbf{Z}_M$ *orbifolds.*

The group generator $\theta \in \Gamma$ acts as follows on the complex coordinates of T^6

$$\theta: \ (z^1, z^2, z^3) \longrightarrow (e^{2\pi i \ v^1} \ z^1 \ , \ e^{2\pi i \ v^2} \ z^2 \ , \ e^{2\pi i \ v^3} \ z^3) \ ,$$

with $\pm v^1 \pm v^2 \pm v^3 = 0$ to furnish $SU(3)$ holonomy ($\theta \ \Omega = \Omega$) [48]. In Table 3, we give a list of possible $\mathbf{Z}_N$ and $\mathbf{Z}_N \times \mathbf{Z}_M$ orbifolds, together with the torus lattices they live on and their Hodge numbers. The lattices marked with $\flat$, $\sharp$, and $*$ are realized as generalized Coxeter twists, the automorphism being in the first and second case $S_1 S_2 S_3 S_4 P_{36} P_{45}$ and in the third $S_1 S_2 S_3 P_{16} P_{25} P_{34}$.

The twist elements $\theta, \ldots, \theta^{N-1}$ produce conical singularities. In a small neighborhood around them, the space locally looks like $\mathbf{C}^3/\Gamma$ (isolated singularity) or $\mathbf{C}^2/\Gamma^{(2)} \times \mathbf{C}$ (non–isolated singularity). In *Ref.* [4] these singularities are resolved using the methods of toric geometry resulting in a smooth Calabi–Yau space Y_6. Afterwards a consistent orientifold action $\mathcal{O}$ is introduced, resulting in the Calabi–Yau orientifold X_6. After resolving the orbifold, three kinds of divisors $\mathcal{D}$ appear, namely E_α, $D_{i\alpha}$, and R_i. The divisors E_α are the exceptional divisors arising from the resolution of an orbifold singularity f_α (or an orbit under the orbifold group), while the divisors $D_{i\alpha}$ denote hyperplanes passsing through fixed points: $D_{i\alpha} = \{z^i = z^i_{fixed,\alpha}\}$. The divisors $R_i = \{z^i = c\}$ for $c \neq z^i_{fixed,\alpha}$ are hyperplanes not passing through a fixed point [4]. As opposed to the $D_{i\alpha}$ they are allowed to move.

As already anticipated in Section 3, some divisors E (or divisor orbits under the orbifold group on the T^6) in the geometry of the covering space Y_6 may not be invariant under the orientifold action σ, *cf. Eq.* (3.7). In this case, a pair of divisors (E_i, E_a), which are eigenstates (with eigenvalues ± 1) under σ may be constructed, *cf. Eq.* (3.9). To this end, the original number of divisors $h_{(1,1)}(Y_6)$ is split into $h^{(+)}_{(1,1)}(X_6)$ even and $h^{(-)}_{(1,1)}(X_6)$ odd divisors. These numbers are determined for the orientifolds of the resolved orbifolds (5.1) in *Ref.* [4] and are displayed in Table 4. In the previous section we have discussed the different stabilization mechanisms for these two kinds of Kähler moduli.

Γ	$h^{(+)}_{(1,1)}$	$h^{(-)}_{(1,1)}$	Γ	$h^{(+)}_{(1,1)}$	$h^{(-)}_{(1,1)}$
$\mathbf{Z}_3$	23	13	$\mathbf{Z}_{8-II}$	27	4
$\mathbf{Z}_4$	25	6	$\mathbf{Z}_{8-II}$	31	0
$\mathbf{Z}_4$	27	4	$\mathbf{Z}_{12-I}$	18	6
$\mathbf{Z}_4$	31	0	$\mathbf{Z}_{12-I}$	22	6
$\mathbf{Z}_{6-I}$	19	6	$\mathbf{Z}_{12-II}$	31	0
$\mathbf{Z}_{6-I}$	23	6	$\mathbf{Z}_2 \times \mathbf{Z}_2$	51	0
$\mathbf{Z}_{6-II}$	19	6	$\mathbf{Z}_2 \times \mathbf{Z}_4$	61	0
$\mathbf{Z}_{6-II}$	23	6	$\mathbf{Z}_2 \times \mathbf{Z}_6$	51	0
$\mathbf{Z}_{6-II}$	21	8	$\mathbf{Z}_2 \times \mathbf{Z}_{6'}$	36	0
$\mathbf{Z}_{6-II}$	25	8	$\mathbf{Z}_3 \times \mathbf{Z}_3$	47	37
$\mathbf{Z}_7$	15	9	$\mathbf{Z}_3 \times \mathbf{Z}_6$	51	22
$\mathbf{Z}_{8-I}$	24	5	$\mathbf{Z}_4 \times \mathbf{Z}_4$	90	0
$\mathbf{Z}_{8-I}$	27	0	$\mathbf{Z}_6 \times \mathbf{Z}_6$	84	0

Table 4: *Hodge numbers $h_{(1,1)}(X_6)$ after the orientifold action*

We choose the orientifold action such that it gives rise to $O3$–planes and $O7$–planes. On the local $\mathbf{C}^3/\Gamma$ patches, an involution, possibly involving the new coordinates associated to the exceptional divisors is chosen, see Section 5 of [4].

Since each $O7$–plane induces -8 units of $D7$–brane charge, we choose to cancel this tadpole locally by placing a stack of 8 coincident $D7$–branes on top of each divisor fixed under the combination of the involution and the scaling action. Each such stack therefore carries an SO(8) gauge group. For the $D3$–brane charge, the case is a bit more involved. The contribution from the $O3$–planes is (in the orientifold quotient X_6 of Y_6)

$$Q_3(O3) = -\frac{1}{4}\, n_{O3}\ ,$$

where n_{O3} denotes the number of $O3$–planes. The $D7$–branes also contribute to the $D3$–tadpole (in the orientifold quotient X_6)

$$Q_3(D7) = -\frac{1}{2}\sum_a \frac{n_{D7,a}\,\chi(\mathcal{D}_a)}{24}\ ,$$

where $n_{D7,a}$ denotes the number of $D7$–branes in the stack located on the divisor $\mathcal{D}_a$. As we have seen, the $\mathcal{D}_a$ can be local D–divisors as well as exceptional divisors E. The last contribution to the $D3$–brane tadpole comes from the $O7$–planes (in the orientifold quotient X_6):

$$Q_3(O7) = -\frac{1}{2}\sum_a \frac{\chi(\mathcal{D}_a)}{6}\ .$$

So the total $D3$–brane charge that must be cancelled is:

$$Q_{3,tot} = -\frac{n_{O3}}{4} - \frac{1}{2} \sum_a \frac{(n_{D7,a}+4)\,\chi(\mathcal{D}_a)}{24} \ . \tag{5.2}$$

These are the values for the orientifold quotient X_6, in the double cover Y_6 this value must be multiplied by two (*cf.* Subsection 5.3). Because we would like to avoid mobile $D3$–branes, this tadpole will be saturated by 3–form flux G_3.

The formula (5.2) for the total $D3$–brane charge $Q_{3,tot}$ differs from the known tadpole equation for the singular orbifold case by the second term. The latter is induced by the curvature of the $D7$–branes which is absent in the singular case. In that case, the number of orientifold $O3$–planes is always 64, *i.e.* $n_{O3} = 64$, and (5.2) boils down to $Q_{3,tot} = -16$ [49]. In the CFT description, this tadpole originates from the total leading divergent contribution of the Klein bottle amplitude $Z_{\mathcal{K}}(1,1)$ of the untwisted orbifold sector. However, there are additional tadpole contributions from other orbifold sectors to be cancelled. More precisely, the tadpole arising from the Klein bottle amplitude $Z_{\mathcal{K}}(1,\theta^k)$ and in addition for even N the $\mathbf{Z}_2$–twisted tadpole related to $Z_{\mathcal{K}}(\theta^{N/2},\theta^k)$ have to be cancelled ($k = 0,\ldots,N-1$). The tadpoles from the sector $(1,1)$ and for even N also from the sector $(1,\theta^{N/2})$ may be cancelled by introducing the right amount of $D3$–brane (or/and 3–form flux) and $D7$–branes, respectively. On the other hand, the divergences of the Klein bottle amplitude $Z_{\mathcal{K}}(1,\theta^k)$, $k \neq 0$ or for even N from the combination $Z_{\mathcal{K}}(1,\theta^k) + Z_{\mathcal{K}}(\theta^{N/2},\theta^k)$, $k \neq 0, N/2$ can only be cancelled against any of the annulus and Möbius strip contributions in the case that the orbifold group Γ is $\mathbf{Z}_3$, $\mathbf{Z}_{6-I}$, $\mathbf{Z}_{6-II}$, $\mathbf{Z}_7$ or $\mathbf{Z}_{12-I}$ [50] or $\mathbf{Z}_2\times\mathbf{Z}_2, \mathbf{Z}_3\times\mathbf{Z}_3, \mathbf{Z}_6\times\mathbf{Z}_6, \mathbf{Z}_2\times\mathbf{Z}_3, \mathbf{Z}_2\times\mathbf{Z}_6, \mathbf{Z}_2\times\mathbf{Z}_6'$ [51]. Hence singular orbifolds have much more constraining tadpole equations, which are non–trivial to fulfill for all $\mathbf{Z}_N$– and $\mathbf{Z}_N\times\mathbf{Z}_M$ orbifolds. However, if one introduces discrete torsion or vector structure tadpoles from all orbifold sectors may be completely cancelled in all singular orbifold cases [52].

Nevertheless, the orientifolds X_6 constructed geometrically in *Ref.* [4] in the large radius regime from resolved orbifolds Y_6 need not have a CFT counterpart in their orbifold limit, since D–branes (in particular stacks of $D7$ and $O7$–branes) wrapping cycles which vanish in the orbifold limit, give rise to extra non–perturbative states in the orbifold limit.

5.2. Resolved toroidal orientifolds as candidate models for a KKLT scenario

In [16], a toroidal orbifold model, namely type IIB string theory compactified on the orientifold of the resolved $T^6/\mathbf{Z}_2\times\mathbf{Z}_2$, was checked for its suitability as a compactification manifold for the KKLT proposal. Since the F–theory lift of this example is known, Witten's criterion could be checked directly and the results of [16] strongly indicate that in this model, all geometric moduli can be fixed.

The methods to obtain a smooth Calabi–Yau manifold from a toroidal orbifold and to subsequently pass to the corresponding orientifold as described in *Ref.* [4] enable us to explicitly check other toroidal orbifolds for their suitability as candidate models for the KKLT proposal.

The requirement that the scalar mass matrix be positive, discussed in Section 2.3, places severe constraints on the list of possible models. Those orbifolds without complex structure moduli do not give rise to stable vacua after the uplift to dS space. Thus $\mathbf{Z}_3$, $\mathbf{Z}_7$, $\mathbf{Z}_{8-I}$ on $SU(4)^2$, $\mathbf{Z}_2 \times \mathbf{Z}_{6'}$, $\mathbf{Z}_3 \times \mathbf{Z}_3$, $\mathbf{Z}_4 \times \mathbf{Z}_4$ and $\mathbf{Z}_6 \times \mathbf{Z}_6$ are excluded from the list of possible models given in Table 3. For more details *cf.* Subsection 5.5.

Since the stabilization of twisted complex structure moduli via 3–form flux is not well understood yet, the models with $h^{twist.}_{(2,1)}(X_6) \neq 0$ cannot be checked explicitly. Yet considerations regarding the topology of their divisors suggest that they might not be suitable candidate models anyway, as will be explained later on. The only models which are not yet excluded and are directly amenable to our methods are thus $T^6/\mathbf{Z}_4$ on $SU(4)^2$, $T^6/\mathbf{Z}_{6-II}$ on $SU(2) \times SU(6)$, the above mentioned $T^6/\mathbf{Z}_2 \times \mathbf{Z}_2$, and $T^6/\mathbf{Z}_2 \times \mathbf{Z}_4$. The example $T^6/\mathbf{Z}_4$ on $SU(4)^2$ contains five instead of the usual three untwisted Kähler moduli. Since it is not clear how these two extra non–diagonal untwisted Kähler moduli contribute to the superpotential, this example will not be discussed explicitly.

The question one would like to answer is: Do enough of the divisors of the above models contribute to the non-perturbative superpotential that all Kähler moduli can be fixed? To answer this question, the topologies of the divisors must be studied. In Section 4.3 of [4] it was shown that there are four basic topologies for the divisors of the resolved toroidal orbifolds: The divisors R_i inherited from the covering T^6 have the topology of either **(i)** $K3$ or **(ii)** T^4. The exceptional divisors E_i which arise in the blowing up process can be birationally equivalent to either **(iii)** a rational surface (*i.e.* $\mathbf{P}^2$ or $\mathbf{F}_n$) or **(iv)** $\mathbf{P}^1 \times T^2$. The same is true for the D–divisors, which correspond to planes fixed at the loci of the fixed points and are linear combinations of the Rs and Es. The rational surfaces have $h_{(1,0)} = h_{(2,0)} = 0$ and therefore $\chi(\mathcal{O}_S) = 1$. Since $h_{(1,0)}$ and $h_{(2,0)}$ are birational invariants, the number of blow–ups which depends on the triangulation of the resolution is irrelevant here. $\mathbf{P}^1 \times T^2$ has $h_{(1,0)} = 1,\ h_{(2,0)} = 0$, T^4 has $h_{(1,0)} = 2,\ h_{(2,0)} = 1$, which both results in $\chi(\mathcal{O}_S) = 0$. $K3$ has $h_{(1,0)} = 0,\ h_{(2,0)} = 1$ and therefore $\chi(\mathcal{O}_S) = 2$.

Since except for $T^6/\mathbf{Z}_2 \times \mathbf{Z}_2$, the F-theory lifts of these models are not known, it must be determined directly in type IIB which divisors contribute to the non–perturbative superpotential. Here, we make use of the index for the Dirac operator on the world–volume of the Euclidean $D3$–brane (4.4). The values of the index for the four divisor topologies arising from resolutions of toroidal orbifolds are given in Table 5.

Topology	$O7$ on top	inters. in 1 dim.	no intersection
$K3$	$2/[1]$	0	$0/[-1]$
T^4	$0/[-1]$	$0/[-2]$	$0/[-3]$
$\mathbf{P}^1 \times T^2$	0	$1/[0]$	$0/[-1]$
$\mathbf{P}^2,\ \mathbf{F}_n$	1	1	0

Table 5: *Index χ_{D3} for the four basic topologies*

The numbers in square brackets are the values of the index in the case that the corresponding zero modes have been lifted by flux, *cf.* Table 2. We see thus that for the case that the $O7$–plane does not intersect the divisor, we never get a contribution, so we better seek an orientifold action which leads to many $O7$–plane solutions. $K3$ can contribute for the case that the $O7$ lies on top of the divisor if the $h^{(+)}_{(2,0)}$ zero modes are lifted by flux. In our set–up, the case that the $O7$ lies on top of the divisor cannot arise, since only the inherited divisors R_i can have the topology of $K3$, and these divisors are never wrapped by $O7$–planes. A divisor with the topology of T^4 can likewise never contribute. $\mathbf{P}^1 \times T^2$ can contribute in case of an intersection with the $O7$–plane in one direction if *no* zero modes are lifted by flux. The rational surfaces always contribute except if there is no intersection irrespective of the background flux. To summarize: All those models are likely to allow the stabilization of all geometric moduli for which

(i) the fixed points and fixed lines are all in equivalence classes with only one member, giving rise to E and D divisors which are birationally equivalent to rational surfaces and

(ii) an orientifold action exists which gives rise to enough $O7$–plane solutions that each divisor intersects an $O7$–plane in at least one complex dimension.

When these conditions are met, it is likely that all geometric moduli will be stabilized when the full scalar potential is minimized.

Requirements (i) and (ii) are both met by $T^6/\mathbf{Z}_4$ on $SU(4)^2$, $T^6/\mathbf{Z}_{6-II}$ on $SU(2) \times SU(6)$, $T^6/\mathbf{Z}_2 \times \mathbf{Z}_2$ and $T^6/\mathbf{Z}_2 \times \mathbf{Z}_4$, therefore we expect that all geometric moduli can be stabilized in these cases.

Models with fixed lines without fixed points on them which lie in orbits of length greater than one do not satisfy criterion (i) since the divisors corresponding to these fixed lines have the topology of $\mathbf{P}^1 \times T^2$. These are exactly the models with $h^{(2,1)}_{twist.} \neq 0$. Unless an elaborate configuration of O–planes can be chosen such that all these divisors intersect on $O7$–plane along one dimension, these examples in general allow only for a partial stabilization of the geometric moduli via Euclidean $D3$–brane instantons. It should be stressed that examples like these are still not completely hopeless since additional effects might lead to the complete stabilization of all moduli. On the other hand, this survey again confirms the old suspicion that manifolds with the right geometrical properties to allow the stabilization of all Kähler moduli by Euclidean $D3$–brane instantons or gaugino condensates are not very generic.

So far, we discussed the conditions for a contribution to the non–perturbative superpotential from Euclidean $D3$–instantons. Since we cancel the $O7$–tadpole by placing $D7$–branes on top of the $O7$–planes, a gaugino condensate can arise on the world–volume of the $D7$–branes. As mentioned before, for a contribution to the non–perturbative superpotential to arise from a gaugino condensate, we should have
(a) no bifundamental matter. This is given when the different divisors on which $D7$–branes are wrapped do not intersect. This condition can be easily checked by inspection of the toric diagram of the resolved patches.
(b) no adjoint matter. This depends on the Hodge numbers of the divisor which is wrapped by the brane. For rational surfaces, *i.e.* $h_{(1,0)} = h_{(2,0)} = 0$, this criterion is fulfilled.

In the following, moduli stabilization will be discussed in detail for the two examples $T^6/\mathbf{Z}_{6-II}$ on $SU(2) \times SU(6)$ and $T^6/\mathbf{Z}_2 \times \mathbf{Z}_4$ on $SU(2)^2 \times SO(5)^2$. In Subsection 5.3 stabilization of the dilaton and complex structure moduli through 3–form flux G_3 is discussed and in Subsection 5.4, the stabilization of the Kähler moduli.

5.3. Complex structure and dilaton stabilization through 3*–form flux*

For the orbifolds X_6 with $h^{(-)}_{(2,1)}(X_6) = 1$ the Kähler potential for the dilaton and complex structure modulus ($U \equiv U^3$) (2.9) is:

$$K_0 = -\log(S + \bar{S}) - \log(U + \bar{U}) \ , \tag{5.3}$$

while the tree–level superpotential (2.14) may be written as

$$W_0 = A + B\ S + U\ (C + D\ S) \ , \tag{5.4}$$

with $A, B, C, D \in \mathbf{C}$ to be specified later. With the F–terms

$$\begin{aligned} \overline{F}^{\overline{S}} &= \left(\frac{S+\overline{S}}{U+\overline{U}}\right)^{1/2} \ [\ -A + B\ \overline{S} - U\ (C - D\ \overline{S})\ ]\ , \\ \overline{F}^{\overline{U}} &= \left(\frac{U+\overline{U}}{S+\overline{S}}\right)^{1/2} \ [\ -A - B\ S + \overline{U}\ (C + D\ S)\ ]\ , \end{aligned} \tag{5.5}$$

we may cast the scalar potential

$$V = g_{S\overline{S}}\ F^S \overline{F}^{\overline{S}} + g_{U\overline{U}}\ F^U \overline{F}^{\overline{U}} - 3\ e^{K_0}\ |W_0|^2$$

into the form:

$$V = \frac{1}{U+\bar{U}}\frac{1}{S+\bar{S}}\ \Big[\ |\ A - B\ \overline{S} + U\ (C - D\ \overline{S})\ |^2 + |\ A + B\ S - \overline{U}\ (C + D\ S)\ |^2 \\ -3\ |\ A + B\ S + U\ (C + D\ S)\ |^2\ \Big]\ . \tag{5.6}$$

The extremal points in the moduli space (S, U) are determined by the solutions of the equations $F^S, F^U = 0$:

$$s_2 = \frac{i}{2}\ \frac{\overline{B}\ C - B\ \overline{C} - \overline{A}\ D + A\ \overline{D}}{\overline{B}\ D + B\ \overline{D}}\quad ,\quad u_2 = \frac{i}{2}\ \frac{-\overline{B}\ C + B\ \overline{C} - \overline{A}\ D + A\ \overline{D}}{\overline{C}\ D + C\ \overline{D}}\ , \tag{5.7}$$

and similarly for the real parts s_1, u_1.

The 3–form flux $G_3 = F_3 + i\ S\ H_3$

$$\frac{1}{(2\pi)^2\alpha'}\ G_3 = \sum_{i=0}^{3}\big[\ (a^i + i\ S\ c^i)\ \alpha_i + (b_i + i\ S\ d_i)\ \beta^i\ \big] + \sum_{j=1}^{6}\big[(e^j + i\ S\ g^j)\ \gamma_j + (f_j + i\ S\ h_j)\ \delta^j\big] \tag{5.8}$$

entering (2.14) is given as linear combination w.r.t. the integer cohomology basis $\{\alpha_i, \beta^i\}_{i=0,\ldots,3}$ and $\{\gamma_j, \delta^j\}_{j=1,\ldots,6}$ [3]. This gives rise to 20 real flux components to be constrained by the respective orbifold group $\mathbf{Z}_N$. This allows to express the complex parameters A, B, C, D through the eight integers $a^0, a^1, b_0, b_1, c^0, c^1, d_0, d_1$. For more details *cf.* [3]. The F–flatness conditions $F^S, F^U = 0$ force the complex structure to align such, that the flux G_3

$$\begin{aligned}\frac{1}{(2\pi)^2\alpha'}\ G_3 = \frac{i}{2\ \mathrm{Re}(U)}\ \Big\{\ &\big[\ \overline{A} - \overline{B}\ S + \overline{U}\ (\overline{C} - \overline{D}\ S)\ \big]\ dz^1 \wedge dz^2 \wedge dz^3 \\ &- \big[\ A + B\ S + U\ (C + D\ S)\ \big]\ d\overline{z}^1 \wedge d\overline{z}^2 \wedge d\overline{z}^3 \\ &+ \big[\ A + B\ S - \overline{U}\ (C + D\ S)\ \big]\ d\overline{z}^1 \wedge d\overline{z}^2 \wedge dz^3 \\ &- \big[\ \overline{A} - \overline{B}\ S - U\ (\overline{C} - \overline{D}\ S)\ \big]\ dz^1 \wedge dz^2 \wedge d\overline{z}^3\ \Big\}\end{aligned} \tag{5.9}$$

becomes ISD, *i.e.* it has only $(2,1)$ and $(0,3)$–components at the extremum. The flux G_3 induces the contribution of

$$N_{flux} = \frac{1}{(2\pi)^4\alpha'^2}\ \int_{Y_6} F_3 \wedge H_3 \tag{5.10}$$

to the total $D3$–brane charge (5.2). Generically, this integral is calculated in the orientifold cover Y_6. Therefore the number N_{flux} has to be twice the negative value of the total $D3$–brane charge (5.2), *i.e.*

$$N_{flux} = -2\ Q_{3,tot} \tag{5.11}$$

to cancel the latter by flux only.

(i) $\mathbf{Z}_{6-II}$ – orbifold on the $SU(2)\times SU(6)$ lattice :

The $\mathbf{Z}_{6-II}$–orbifold has the action $(v^1,\, v^2,\, v^3) = (\frac{1}{6},\, \frac{1}{3},\, -\frac{1}{2})$. The 3–form flux (5.8) constrained by the $\mathbf{Z}_{6-II}$–orbifold group becomes:

$$\begin{aligned} \frac{1}{(2\pi)^2\alpha'}\; G_3 =& \frac{1}{3}\; (a_0+iSc_0)\; (3\;\alpha_0+2\;\beta_3+\gamma_1-2\gamma_2-2\;\gamma_3+\gamma_4-\delta_5) \\ & +(b_0+iSd_0)\;(-\alpha_3+\beta_0+\gamma_5-\gamma_6) \\ & +\frac{1}{2}\;(b_1+iSd_1)\;(2\;\beta_1+\beta_2+\delta_1-\delta_2-2\;\delta_3-\delta_4) \\ & +(a_1+iSc_1)\;(\alpha_1+\alpha_2+\beta_3-\gamma_2-\gamma_3-\delta_6)\ . \end{aligned} \tag{5.12}$$

This flux correspond to the flux number:

$$N_{\text{flux}} = 2\; b_0\; c_0 + b_1\; (c_0+3\; c_1) - 2\; a_0\; d_0 - d_1\; (a_0+3\; a_1)\ . \tag{5.13}$$

For the $\mathbf{Z}_{6-II}$ orbifold with $SU(2)\times SU(6)$ lattice the coefficients A, B, C, D entering (5.6) become:

$$\begin{aligned} &A = -\frac{\sqrt{3}}{2}\; b_1 + ib_0 \quad , \quad B = -d_0 - \frac{\sqrt{3}\,i}{2} d_1\ , \\ &C = a_0 + i\left(\frac{a_0}{\sqrt{3}}+\sqrt{3}a_1\right) \quad , \quad D = -\left(\frac{c_0}{\sqrt{3}}+\sqrt{3}c_1\right)+ic_0\ . \end{aligned} \tag{5.14}$$

With this information, *Eq.* (5.4) yields the superpotential:

$$\begin{aligned} W_0 =& -\frac{\sqrt{3}}{2}\; b_1 + i\; b_0 - S\; \left(d_0+\frac{\sqrt{3}i}{2}\; d_1\right) \\ & +U\;\left[\;a_0+i\left(\frac{a_0}{\sqrt{3}}+\sqrt{3}\;a_1\right)\;\right] - S\;U\;\left(\frac{c_0}{\sqrt{3}}+\sqrt{3}\;c_1-i\;c_0\right)\ . \end{aligned} \tag{5.15}$$

Since the total $D3$–brane charge in the CY orientifold is $Q_{3,tot}=-22$ (see Section 5.4), we look for fluxes (5.12) with $N_{flux}=44$ on the covering space Y_6. Furthermore, the fields $S=s_1+is_2$ and $U=u_1+iu_2$ should be fixed (*cf.* (5.7)) to realistic values. A reasonable value for ReS is $s_1\sim 3.6$, which corresponds to a string coupling constant $g_{\text{string}}\sim 0.27$ at the string scale. Besides, the complex structure modulus U is expected to be around the ρ–point in the fundamental region, with $\rho=\frac{1}{2}+\frac{i}{2}\sqrt{3}$. An additional constraint may be imposed on the tuning parameter $e^{K_0}|W_0|^2$, which should be small to avoid higher order effects in the full non–perturbative superpotential (2.13). After a systematic scan in the flux space $(a^0, a^1, b_0, b_1, c^0, c^1, d_0, d_1)\in\mathbf{Z}^8$ we find hundreds of vacua, which meet these criteria. A set of equivalent vacua, differing only in the discrete flux parameters $(a^0, a^1, b_0, b_1, c^0, c^1, d_0, d_1)$, is given in the following Table 6.

$(a^0,\ b_0,\ c^0,\ d_0,\ a^1,\ b_1,\ c^1,\ d_1)$	s_1	s_2	u_1	u_2	m_S	m_U
$(-5,\ 12,\ 0,\ 2,\ -4,\ -8,\ -1,\ 0)$	3.15788	5.83333	1.26315	0.0666667	2.18	13.68
$(-5,\ 10,\ 0,\ 2,\ -3,\ -8,\ -1,\ 0)$	3.15788	4.83333	1.26315	0.0666667	2.18	13.68
$(-5,\ 6,\ 0,\ 2,\ -1,\ -8,\ -1,\ 0)$	3.15788	2.83333	1.26315	0.0666667	2.18	13.68
$(-5,\ 0,\ 0,\ 2,\ 2,\ -8,\ -1,\ 0)$	3.15788	−0.166667	1.26315	0.0666667	2.18	13.68
$(-5,\ -4,\ 0,\ 2,\ 4,\ -8,\ -1,\ 0)$	3.15788	−2.16667	1.26315	0.0666667	2.18	13.68
$(-5,\ -8,\ 0,\ 2,\ 6,\ -8,\ -1,\ 0)$	3.15788	−4.16667	1.26315	0.0666667	2.18	13.68
$(-5,\ -12,\ 0,\ 2,\ 8,\ -8,\ -1,\ 0)$	3.15788	−6.16667	1.26315	0.0666667	2.18	13.68
$(5,\ 10,\ 0,\ -2,\ -7,\ 8,\ 1,\ 0)$	3.15788	−5.16667	1.26315	0.0666667	2.18	13.68
$(5,\ 8,\ 0,\ -2,\ -6,\ 8,\ 1,\ 0)$	3.15788	−4.16667	1.26315	0.0666667	2.18	13.68
$(5,\ 6,\ 0,\ -2,\ -5,\ 8,\ 1,\ 0)$	3.15788	−3.16667	1.26315	0.0666667	2.18	13.68
$(5,\ 2,\ 0,\ -2,\ -3,\ 8,\ 1,\ 0)$	3.15788	−1.16667	1.26315	0.0666667	2.18	13.68

Table 6: *Discrete landscape of supersymmetric AdS minima for* $N_{flux} = 44$, $e^{K_0/2}\ |W_0| = 0.34864$ and $V_0 = -0.364644$.

Clearly, the axionic vev s_2 may be shifted back into the fundamental region $s_2 \equiv\ -0.166667$, while the flux number N_{flux} in (5.10) and K_0, W_0 are preserved [53]. Furthermore, in Table 7 we present a set of supersymmetric AdS minima in the (S, U)–space with different tuning parameters $e^{K_0}|W_0|^2$.

$(a^0,\ b_0,\ c^0,\ d_0,\ a^1,\ b_1,\ c^1,\ d_1)$	s_1	s_2	u_1	u_2	$-V_0$	$e^{K/2}\ \|W_0\|$	m_S	m_U
$(\ -2,\ -7,\ -1,\ 2,\ 3,\ -8,\ 0,\ -2)$	3.8092	−0.3	2.11622	0.722222	0.02273	0.087046	1.52	4.91
$(\ 0,\ -10,\ -1,\ -1,\ 5,\ -3,\ 1,\ -2)$	3.7944	3.95	1.2648	0.316667	0.278999	0.304958	1.51	13.67
$(\ 0,\ -10,\ -1,\ 2,\ 3,\ -6,\ 0,\ -2)$	3.7934	−1.9	2.10745	0.944444	0.296119	0.314176	1.51	4.92
$(\ 2,\ -10,\ -1,\ -1,\ 3,\ -1,\ 1,\ -2)$	3.7918	2.65	1.26392	0.55	0.324662	0.328969	1.51	13.67
$(\ 0,\ -10,\ -1,\ -1,\ 6,\ -6,\ 1,\ -2)$	3.7296	4.7	1.036	0.305556	1.40124	0.683432	1.51	19.86
$(\ 0,\ -10,\ -1,\ -1,\ 4,\ 0,\ 1,\ -2)$	3.7095	3.2	1.5456	0.333333	1.75049	0.763869	1.51	8.87
$(\ 5,\ -9,\ -1,\ -1,\ 0,\ 3,\ 1,\ -2)$	3.6575	0.35	1.21918	0.783333	2.64978	0.93982	1.51	13.95

Table 7: *Supersymmetric AdS minima in the* (S, U)*–space for* $N_{flux} = 44$ *and specific* $e^{K_0/2}\ |W_0|$.

$(ii)\ \ \mathbf{Z}_2 \times \mathbf{Z}_4$ – orbifold on the $SU(2)^2 \times SO(5)^2$ lattice :

The $\mathbf{Z}_2 \times \mathbf{Z}_4$–orbifold has the two actions $(v^1, v^2, v^3) = \frac{1}{2}(1, 0, -1)$ and $(w^1, w^2, w^3) =$

$\frac{1}{4}(0,1,-1)$. The 3–form flux (5.8) constrained by the $\mathbf{Z}_2 \times \mathbf{Z}_4$–orbifold group becomes:

$$\begin{aligned}\frac{1}{(2\pi)^2\alpha'}\, G_3 =& (a_3 + iSc_0)\,(-\alpha_2 + \alpha_3) + (a_0 + iSc_0)\,\left(\alpha_0 - \alpha_2 - \frac{1}{2}\beta_1\right) \\ &+ (b_2 + iSd_2)\,\left(\alpha_1 + \frac{1}{2}\beta_0 + \beta_2\right) + (b_3 + iSd_3)\,\left(\alpha_1 + \frac{1}{2}\beta_0 + \beta_3\right)\ .\end{aligned} \tag{5.16}$$

The coefficients A, B, C, D entering (5.6) are given in the case of $\mathbf{Z}_2 \times \mathbf{Z}_4$–orbifold with $SU(2)^2 \times SO(5)^2$ lattice by:

$$\begin{aligned} A &= -\frac{1+2}{2}\,(b_2 - i\, b_3) \quad , \quad B = \frac{1-i}{2}\,(d_2 - i\, d_3)\ , \\ C &= \frac{1+i}{2}a_0 + a_3 \quad , \quad D = \frac{-1+i}{2}c_0 + i\, c_3\ . \end{aligned} \tag{5.17}$$

Furthermore, the flux number is:

$$N_{\rm flux} = a_3 d_2 - b_2 c_3 + b_3\,(c_0 + c_3) - (a_0 + a_3)\,d_3\ . \tag{5.18}$$

With this information, the superpotential (5.4) becomes:

$$\begin{aligned} W_0 =& -\frac{1+i}{2}\,(b_2 - i\, b_3) + S\,\frac{(1-i)}{2}\,(d_2 - i\, d_3) \\ &+ U\left(\frac{1+i}{2}\,a_0 + a_3\right) + S\,U\,\left(\frac{-1+i}{2}\,c_0 + i\, c_3\right)\ . \end{aligned} \tag{5.19}$$

We search for fluxes (5.16) with $N_{flux} = 52$. We fix the value of the s_1 at 3.24, which corresponds to a string coupling constant $g_{\rm string} = 0.30$ at the string scale. A set of equivalent vacua, differing only in the discrete flux parameters $(a^0, b_2, c^0, d_2, a^3, b_3, c^3, d_3)$, is given in the following Table 8.

$(a^0,\ b_2,\ c^0,\ d_2,\ a^3,\ b_3,\ c^3,\ d_3)$	s_1	s_2	u_1	u_2	m_S	m_U
$(14,\ 14,\ 0,\ 1,\ -3,\ 14,\ -2,\ -5)$	3.23796	−1.875	0.925131	−1.46429	2.46	30.21
$(14,\ 15,\ 0,\ 1,\ -3,\ 15,\ -2,\ -5)$	3.23796	−2.125	0.925131	−1.53571	2.46	30.21
$(-8,\ 8,\ 4,\ 1,\ 11,\ 8,\ -2,\ -3)$	3.23796	−1.875	0.925131	0.535714	2.46	30.21
$(-14,\ 14,\ 0,\ 5,\ 11,\ 14,\ 2,\ -1)$	3.23796	1.875	0.925131	1.46429	2.46	30.21
$(-14,\ 15,\ 0,\ 5,\ 11,\ 15,\ 2,\ -1)$	3.23796	2.125	0.925131	1.53571	2.46	30.21
$(8,\ 8,\ 4,\ 3,\ 3,\ 8,\ -2,\ -1)$	3.23796	1.875	0.925131	−0.535714	2.46	30.21
$(14,\ -16,\ 0,\ -3,\ -19,\ -16,\ -2,\ -1)$	3.23796	6.125	0.925131	0.535714	2.46	30.21
$(14,\ -15,\ 0,\ -3,\ -19,\ -15,\ -2,\ -1)$	3.23796	5.875	0.925131	0.464286	2.46	30.21
$(14,\ -14,\ 0,\ -5,\ -11,\ -14,\ -2,\ 1)$	3.23796	1.875	0.925131	1.46429	2.46	30.21

Table 8: *Discrete landscape of supersymmetric AdS minima for* $N_{flux} = 52$, $e^{K_0/2}\ |W_0| = 0.310374$ and $V_0 = -0.288997$.

In the next table, we present a set of supersymmetric AdS minima in the (S, U)-space with same tuning parameter $e^{K_0}|W_0|^2$, but different choices for S and U.

$(a^0,\ b_2,\ c^0,\ d_2,\ a^3,\ b_3,\ c^3,\ d_3)$	s_1	s_2	u_1	u_2	m_S	m_U
$(-13,\ 17,\ 3,\ 1,\ 11,\ 17,\ -1,\ -5)$	3.70	−2.85714	1.29518	1.85	1.88	15.42
$(-10,\ 16,\ 3,\ 1,\ 9,\ 16,\ -1,\ -5)$	3.70	−2.14286	1.52375	1.82353	1.88	11.14
$(9,\ 15,\ 1,\ 3,\ -1,\ 15,\ -2,\ -5)$	3.70	−1.14286	1.52375	−2.17647	1.88	11.14
$(-14,\ 20,\ 3,\ -1,\ 16,\ 20,\ -2,\ -4)$	3.70	−6.14286	1.29518	1.15	1.88	15.41
$(-9,\ 15,\ 1,\ 5,\ 8,\ 15,\ 1,\ -3)$	3.70	1.14286	1.52375	2.17647	1.88	11.14
$(-4,\ 15,\ 2,\ 2,\ 6,\ 15,\ -1,\ -5)$	3.70	−1.85714	3.23796	1.625	1.87	2.48
$(2,\ 20,\ 1,\ -1,\ 13,\ 20,\ -2,\ -3)$	3.70	−8.14286	1.52375	−0.176471	1.88	11.14
$(-6,\ 20,\ 2,\ -1,\ 12,\ 20,\ -2,\ -4)$	3.24	−5.875	2.15864	0.583333	2.46	5.56
$(-5,\ 10,\ 3,\ 2,\ 7,\ 10,\ -1,\ -4)$	3.24	−0.875	1.61898	1.0625	2.46	9.87
$(3,\ 11,\ 2,\ 2,\ 3,\ 11,\ -2,\ -4)$	3.24	−1.375	2.15864	−0.916667	2.46	5.56
$(-2,\ 15,\ 2,\ 3,\ 4,\ 15,\ -1,\ -5)$	3.24	−0.875	4.31728	1.16667	1.38	2.47
$(-6,\ 14,\ 2,\ 4,\ 6,\ 14,\ 0,\ -5)$	3.24	−0.125	2.15864	2.41667	2.46	5.56

Table 9: *Supersymmetric AdS minima in the* (S, U)*–space for* $N_{flux} = 52$, $e^{K_0/2}\ |W_0| = 0.310374$ *and* $V_0 = -0.28900$.

(*iii*) $\mathbf{Z}_4$ – orbifold on the $SU(4)^2$ lattice :

The $\mathbf{Z}_4$–orbifold has the action $(v^1,\ v^2,\ v^3) = (\frac{1}{4},\ \frac{1}{4},\ -\frac{1}{2})$. The 3–form flux (5.8) constrained by the $\mathbf{Z}_4$–orbifold group becomes

$$\begin{aligned}\frac{1}{(2\pi)^2\alpha'}G_3 =&(a_0 + iSc_0)\ (\alpha_0 + \alpha_3 + \beta_2 - \gamma_2 - \gamma_3 - \delta_5)\\ &+ \frac{1}{2}\ (b_0 + iSd_0)\ (-\alpha_2 + 2\beta_0 + \beta_3 + \gamma_4 + \gamma_5 - \gamma_6 + \delta_3)\\ &+ \frac{1}{2}\ (b_1 + iSd_1)\ (\alpha_2 + 2\beta_1 + \beta_3 - \gamma_4 + \gamma_5 - \gamma_6 - 2\delta_2 - \delta_3)\\ &+ (a_1 + iSc_1)\ (\alpha_1 + 2\alpha_3 + \beta_2 - \gamma_2 - 2\gamma_3 + \delta_4 - \delta_5 - \delta_6)\ .\end{aligned} \tag{5.20}$$

In the case of the $\mathbf{Z}_4$–orbifold with the lattice $SU(4)^2$ the coefficients A, B, C, D entering (5.6) are given by:

$$\begin{aligned}&A = -b_1 + i\ b_0 \quad , \quad B = -d_0 - i\ d_1\ ,\\ &C = a_0 + i\ (a_0 + 2a_1) \quad , \quad D = -c_0 - 2c_1 + i\ c_0\ .\end{aligned} \tag{5.21}$$

Furthermore, the flux number is:

$$N_{\text{flux}} = 2\ [\ (b_0+b_1)\ c_0 + b_1 c_1 - 2a_1 d_1 - (d_0+d_1)\ a_0\ ]\ . \tag{5.22}$$

With this information the superpotential (5.4) becomes:

$$\begin{aligned} W_0 = & -b_1 + i\ b_0 - S\ \ (d_1 + i\ d_0) \\ & + U\ \ [\ a_0 + i\ (a_0 + 2\ a_1)\ ] - S\ U\ \ (c_0 + 2\ c_1 - i\ c_0)\ . \end{aligned} \tag{5.23}$$

5.4. Kähler moduli stabilization

We consider the racetrack superpotential (2.13)

$$W = \widetilde{W}_0 + \sum_{j=1}^{h^{(+)}_{(1,1)}(X_6)} \gamma_j\ e^{a_j\ T^j}\ , \tag{5.24}$$

with $\widetilde{W}_0$ related to the tree–level flux superpotential (5.4), by $\widetilde{W}_0 = -e^{K_0/2}|W_0|$. The redefined quantity $\widetilde{W}_0$ makes sure, that the minimization procedure w.r.t. the set of Kähler moduli $T^1,\ldots,T^n$ in the Kähler gauge $K_0 \equiv 0$ yields the correct value $-3e^K|\widetilde{W}_0|^2 = -3e^{K_0+K}|W_0|^2$ in the scalar potential (2.22). This value accounts for the contribution of the dilaton and complex structure stabilization procedure, which is decoupled and performed in the previous subsection. Here and in the following K is the Kähler potential (2.12) for the $n = h^{(+)}_{(1,1)}(X_6)$ Kähler moduli $T^1,\ldots,T^n$. According to Subsection 2.2, we may assume $\gamma_j \in \mathbf{R}^+$, *i.e.* any complex phase of γ_j has been put into the axionic vevs (2.19) of the Kähler moduli T^j. The supersymmetric vacua are given by the equations (2.17), *i.e.* by the critical points of $e^{K/2}W$. These equations fix the real part of the Kähler moduli T^j, *i.e.* the divisor volumes $\text{Vol}(\mathcal{D}_i)$ of an even divisor $\mathcal{D}_j$:

$$\text{Vol}(\mathcal{D}_j) = \text{Re}(T^j) = \frac{3}{4}\ \mathcal{K}_{ijk}\ t^j\ t^k = \frac{3}{2}\ \frac{\partial}{\partial t^j}\ \text{Vol}(X_6)\ . \tag{5.25}$$

To ignore α'–corrections, the Kähler moduli T^j or divisor volumes $\text{Vol}(\mathcal{D}_j)$ should be stabilized at large values, resulting in a large CY volume $\text{Vol}(X_6)$. The F–flatness conditions (2.17) roughly give rise to the relations $\widetilde{W}_0 \sim \gamma_j\ e^{a_j\ T^j}$. Hence, a smaller $\widetilde{W}_0$ or larger coefficients γ_j yield larger divisor volumes $\text{Re}T^j$. Hence a small $\widetilde{W}_0$ or large divisor volumes guarantee that α'–corrections may be neglected. In (5.24), the exponentials $e^{a_j\ T^j}$ should be small $\sim \mathcal{O}(10^{-4})$, such that multi–instanton processes or multi–wrappings may be neglected. In principle, this means that $\widetilde{W}_0$ should be also of this order $\sim 10^{-4}$ [1]. Furthermore, in *Ref.* [1] it has been argued that due to the smallness of these exponentials a dependence of the coefficients γ_j on the dilaton S and complex structure moduli U^j does not change the critical points of the dilaton and complex structure moduli much, as derived in the previous subsection, as long as the relative derivatives $\gamma_j^{-1}\,\partial_{S,U^\lambda}\gamma_j$ and $\gamma_j^{-1}\,\partial^2_{S,U^\lambda}\gamma_j$ are not huge.

(i) $\mathbf{Z}_{6-II}$ – orbifold on the $SU(2)\times SU(6)$ lattice :

We consider the resolved $\mathbf{Z}_{6-II}$ orbifold Y_6 on the lattice $SU(2)\times SU(6)$ which has $h_{(1,1)}(Y_6)=25$.



Fig. 1: *Schematic picture of the fixed set configuration of* $\mathbf{Z}_{6-II}$ *on* $SU(2)\times SU(6)$

The configuration of the fixed point set is displayed in Figure 1 in a schematic way, where each complex coordinate is shown as a coordinate axis and the opposite faces of the resulting cube of length 1 are identified. We see that there are 12 local $\mathbf{C}^3/\mathbf{Z}_{6-II}$ patches which each sit at the intersection of two fixed lines, 3 $\mathbf{C}^2/\mathbf{Z}_3$ fixed lines in the z^3 direction originating from the order three element θ^2 and 4 $\mathbf{C}^2/\mathbf{Z}_2$ fixed lines in the z^2 direction originating from the order two element θ^3.

The resolution of the $\mathbf{C}^3/\mathbf{Z}_{6-II}$ singularity is described by the toric diagram in Figure 2. From these two figures, we can read off the exceptional divisors [4], which together with the inherited divisors R_i form a basis for $H^{(1,1)}(X_6)$:

$$R_1\quad,\quad R_2\quad,\quad R_3\quad,\quad E_{1,\beta\gamma}\quad,\quad E_{3,\gamma}\quad,\quad E_{2,\beta}\quad,\quad E_{4,\beta}\ , \tag{5.26}$$

with $\beta=1,2,3$, $\gamma=1,\ldots,4$. In addition, the orbifold fixed points give rise to the eight divisors $D_1, D_{2,\beta}$ and $D_{3,\gamma}$. The topologies of these divisors were determined in [4]. $E_{1,\beta\gamma}$



Fig. 2: *Toric diagram of the resolution of* $\mathbf{C}^3/\mathbf{Z}_{6-II}$ *and dual graph*

is a blow–up of $\mathbf{F}_1$ in two points, while the remaining exceptional divisors $E_{2,\beta}, E_{3,\gamma}, E_{4,\beta}$ are all $\mathbf{P}^1 \times \mathbf{P}^1$. $D_1, D_{2\beta}, D_{3\gamma}$ are blow–ups of $\mathbf{P}^1 \times \mathbf{P}^1$ in 12, 8, and 9 points, respectively. Finally, R_1 is a T^4 and R_2, R_3 are K3 surfaces.

Any orientifold projection $\mathcal{O}$ with $O3$– and $O7$–planes is such that the twelve divisors $E_{1,2\gamma}$, $E_{1,3\gamma}$, $E_{2,2}$, $E_{2,3}$, $E_{4,2}$, and $E_{4,3}$ are not invariant under the orientifold action σ [4]:

$$\begin{aligned} \sigma\ E_{1,2\gamma} = E_{1,3\gamma} \quad &, \quad \sigma\ E_{1,3\gamma} = E_{1,2\gamma}\ , \\ \sigma\ E_{2,2} = E_{2,3} \quad &, \quad \sigma\ E_{2,3} = E_{2,2}\ , \\ \sigma\ E_{4,2} = E_{4,3} \quad &, \quad \sigma\ E_{4,3} = E_{4,2}\ , \end{aligned} \tag{5.27}$$

and the six pairs of eigenstates $(E, \widetilde{E})$ under σ have to be constructed (*cf.* also Section 3):

$$\begin{aligned} E_{1,\gamma} := \frac{1}{2}\ (\ E_{1,2\gamma} + E_{1,3\gamma}\ ) \quad &, \quad \widetilde{E}_{1,\gamma} := \frac{1}{2}\ (\ E_{1,2\gamma} - E_{1,3\gamma}\ )\ , \\ E_2 := \frac{1}{2}\ (\ E_{2,2} + E_{2,3}\ ) \quad &, \quad \widetilde{E}_2 := \frac{1}{2}\ (\ E_{2,2} - E_{2,3}\ )\ , \\ E_4 := \frac{1}{2}\ (\ E_{4,2} + E_{4,3}\ ) \quad &, \quad \widetilde{E}_4 := \frac{1}{2}\ (\ E_{4,2} - E_{4,3}\ )\ . \end{aligned} \tag{5.28}$$

As a consequence, we have $h^{(-)}_{(1,1)}(X_6) = 6$ (*cf.* also Table 4). Furthermore, the divisors $D_{2,2}$ and $D_{2,3}$ are mapped to each other under σ and only the combination $D_2 = \frac{1}{2}(D_{2,2} + D_{2,3})$ is invariant. To summarize, the orientifold action $\mathcal{O}$ splits the divisors (5.26) into the even divisors

$$H^{(4)}_{+}(X_6)\ \ni\ E_{1,1\gamma}\ ,\ E_{1,\gamma}\ ,\ E_{2,1}\ ,\ E_2\ ,\ E_{4,1}\ ,\ E_4\ ,\ E_{3,\gamma}\ ,\ D_1\ ,\ D_2\ ,\ D_{2,1}\ ,\ D_{3,\gamma}\ ,\ R_1,\ ,\ R_2\ ,\ R_3 \tag{5.29}$$

and into the odd divisors

$$H^{(4)}_{-}(X_6)\ \ni\ \widetilde{E}_2\quad ,\quad \widetilde{E}_4\quad ,\quad \widetilde{E}_{1,\gamma}\quad ,\quad \widetilde{D}_2 \tag{5.30}$$

with $\gamma = 1,\ldots,4$. We choose the orientifold action σ such that its fixed point set consists of seven $O7$–planes wrapped on the divisors $D_1, D_{3,\gamma}, E_{2,1}$ and E_2. In addition, there are twelve $O3$–planes at $z^2 = 0\ ,\ z^1 \neq 0$. Because of $\chi(D_1) = 16$, $\chi(D_{3,\gamma}) = 13$ and $\chi(E_{2,1}) = \chi(E_2) = 4$, the total $D3$–brane charge $Q_{3,tot}$ in *Eq.* (5.2) is $Q_{3,tot} = -22$. The Poincaré dual 2–forms ω_i of the 19 invariant divisors represent a basis for the Kähler form J:

$$\begin{aligned} J = r_1\ R_1 + r_2\ R_2 + r_3\ R_3 - t_2\ E_2 - t_4\ E_4 - t_{2,1}\ E_{2,1} - t_{4,1}\ E_{4,1} \\ -\sum_{\gamma=1}^{4}\left(\ t_{1,\gamma}\ E_{1,\gamma} + t_{1,1\gamma}\ E_{1,1\gamma} + t_{3,\gamma}\ E_{3,\gamma}\ \right)\ , \end{aligned} \tag{5.31}$$

with the 19 Kähler coordinates $r_1,\ r_2,\ r_3,\ t_{1,\gamma},\ t_{1,1\gamma},\ t_2,\ t_{2,1},\ t_{3,\gamma},\ t_4,\ t_{4,1}$. The intersection numbers computed in [4] are split (*cf.* Subsection 2.1) into the two sets of intersection numbers $\mathcal{K}_{ijk}$

$$\begin{aligned} & R_1R_2R_3 = 3\quad ,\quad R_3E_{2,1}E_{4,1} = 1\quad ,\quad R_3E_2E_4 = \frac{1}{2}\quad ,\quad R_2E_{3,\gamma}^2 = -1\ , \\ & R_3E_{2,1}^2 = -4\quad ,\quad R_3E_{4,1}^2 = -1\quad ,\quad R_3E_2^2 = -2\quad ,\quad R_3E_4^2 = -\frac{1}{2}\ , \\ & E_{1,1\gamma}^3 = 3\quad ,\quad E_{1,\gamma}^3 = \frac{3}{4}\quad ,\quad E_{2,1}^2E_{4,1} = -4\quad ,\quad E_2^2E_4 = -1\ , \\ & E_{2,1}^3 = 32\quad ,\quad E_{4,1}^3 = 4\quad ,\quad E_2^3 = 8\quad ,\quad E_4^3 = 1\quad ,\quad E_{3,\gamma}^3 = 4\ , \\ & E_{1,1\gamma}E_{2,1}^2 = -4\quad ,\quad E_{1,1\gamma}E_{4,1}^2 = -1\quad ,\quad E_{1,1\gamma}E_{3,\gamma}^2 = -1\ , \\ & E_{1,\gamma}E_2^2 = -1\quad ,\quad E_{1,\gamma}E_4^2 = -\frac{1}{4}\quad ,\quad E_{1,\gamma}E_{3,\gamma}^2 = -1\ , \\ & E_{1,1\gamma}E_{2,1}E_{4,1} = 1\quad ,\quad E_{1,\gamma}E_2E_4 = \frac{1}{4}\ , \end{aligned} \tag{5.32}$$

and $\mathcal{K}_{iab}$:

$$\begin{aligned} & R_3\widetilde{E}_2\widetilde{E}_4 = \frac{1}{2}\quad ,\quad R_3\widetilde{E}_2^2 = -2\quad ,\quad R_3\widetilde{E}_4^2 = -\frac{1}{2}\ , \\ & E_{1,\gamma}\widetilde{E}_{1,\gamma}^2 = \frac{3}{4}\quad ,\quad E_4\widetilde{E}_2^2 = -1\quad ,\quad E_2\widetilde{E}_2\widetilde{E}_4 = -1\ , \\ & E_2\widetilde{E}_2^2 = 8\quad ,\quad E_4\widetilde{E}_4^2 = 1\ , \\ & \widetilde{E}_{1,\gamma}\widetilde{E}_2E_2 = -1\quad ,\quad E_{1,\gamma}\widetilde{E}_2^2 = -1\quad ,\quad \widetilde{E}_{1,\gamma}\widetilde{E}_4E_4 = -\frac{1}{4}\quad ,\quad E_{1,\gamma}\widetilde{E}_4^2 = -\frac{1}{4}\ , \\ & E_{1,\gamma}\widetilde{E}_2\widetilde{E}_4 = \frac{1}{4}\quad ,\quad \widetilde{E}_{1,\gamma}E_2\widetilde{E}_4 = \frac{1}{4}\quad ,\quad \widetilde{E}_{1,\gamma}\widetilde{E}_2E_4 = \frac{1}{4}\ . \end{aligned} \tag{5.33}$$

The volume $\mathrm{Vol}(X_6)=\frac{1}{6}\ \int_{X_6} J\wedge J\wedge J=\frac{1}{6}\ \mathcal{K}_{ijk}\ t^i\ t^j\ t^k$ of the CY orientifold X_6 becomes:

$$\begin{aligned}\mathrm{Vol}(X_6)=&\,3\ r_1\ r_2\ r_3+r_3\ (t_{2,1}\ t_{4,1}+\frac{1}{2}\ t_2\ t_4)-\sum_{\gamma=1}^{4}\ (\ t_{1,1\gamma}\ t_{2,1}\ t_{4,1}+\frac{1}{4}\ t_{1,\gamma}\ t_2\ t_4\ )\\&-\frac{1}{2}\ r_2\ \sum_{\gamma=1}^{4}\ t_{3,\gamma}^2-r_3\ (\ 2\ t_{2,1}^2+\frac{1}{2}\,t_{4,1}^2+t_2^2+\frac{1}{4}\ t_4^2\ )\\&-\frac{1}{2}\sum_{\gamma=1}^{4}(\ t_{1,1\gamma}^3+\frac{1}{4}\ t_{1,\gamma}^3\ )+2\ t_{2,1}^2\ t_{4,1}+\frac{1}{2}\ t_2^2\ t_4-\frac{4}{3}\ (\ 4\,t_{2,1}^3+\frac{1}{2}\,t_{4,1}^3+t_2^3+\frac{1}{8}\ t_4^3\ )-\frac{2}{3}\ \sum_{\gamma=1}^{4}t_{3,\gamma}^3\\&+\sum_{\gamma=1}^{4}\ \left(\ 2\ t_{1,1\gamma}\ t_{2,1}^2+\frac{1}{2}\ t_{1,1\gamma}\ t_{3,\gamma}^2+\frac{1}{2}\,t_{1,1\gamma}\ t_{4,1}^2+\frac{1}{2}\ t_{1,\gamma}\ t_2^2+\frac{1}{2}\ t_{1,\gamma}\ t_{3,\gamma}^2+\frac{1}{8}\ t_{1,\gamma}\ t_4^2\ \right)\ .\end{aligned}\tag{5.34}$$

According to *Eq.* (5.25) from (5.32) or (5.34), the 19 divisor volumes are derived:

$$\begin{aligned}\mathrm{Vol}(R_1)&=\frac{9}{2}\ r_2\ r_3\quad,\qquad Vol(R_2)=\frac{9}{2}\ r_1\ r_3-\frac{3}{4}\ \sum_{\gamma=1}^{4}t_{3,\gamma}^2\ ,\\ \mathrm{Vol}(R_3)&=\frac{9}{2}\ r_1\ r_2+\frac{3}{2}\ t_{2,1}\ t_{4,1}+\frac{3}{4}\ t_2\ t_4-3\ t_{2,1}^2-\frac{3}{4}\ t_{4,1}^2-\frac{3}{2}\ t_2^2-\frac{3}{8}\ t_4^2\ ,\\ \mathrm{Vol}(E_{1,\gamma})&=-\frac{9}{16}\ t_{1,\gamma}^2+\frac{3}{4}\ t_2^2+\frac{3}{16}\ t_4^2+\frac{3}{4}\ t_{3,\gamma}^2-\frac{3}{8}\ t_2\ t_4\ ,\\ \mathrm{Vol}(E_{1,1\gamma})&=-\frac{9}{4}\ t_{1,1\gamma}^2+3\ t_{2,1}^2+\frac{3}{4}\ t_{4,1}^2+\frac{3}{4}\ t_{3,\gamma}^2-\frac{3}{2}\ t_{2,1}\ t_{4,1}\ ,\\ \mathrm{Vol}(E_2)&=\frac{3}{4}\ r_3\ t_4-3\ r_3\ t_2+\frac{3}{2}\ t_2\ t_4-6\ t_2^2-\frac{3}{8}\ \sum_{\gamma=1}^{4}(\ t_{1,\gamma}\ t_4-4t_2\ t_{1,\gamma}\ )\ ,\\ \mathrm{Vol}(E_{2,1})&=\frac{3}{2}\ r_3\ t_{4,1}-6\ r_3\ t_{2,1}+6\ t_{2,1}\ t_{4,1}-24\ t_{2,1}^2-\frac{3}{2}\ \sum_{\gamma=1}^{4}(\ t_{1,1\gamma}\ t_{4,1}-4t_{2,1}\ t_{1,1\gamma}\ )\ ,\\ \mathrm{Vol}(E_{3,\gamma})&=-\frac{3}{2}\ r_2\ t_{3,\gamma}-3\ t_{3,\gamma}^2+\frac{3}{2}\ t_{1,1\gamma}\ t_{3,\gamma}+\frac{3}{2}\ t_{1,\gamma}\ t_{3\gamma}\ ,\end{aligned}\tag{5.35}$$

$$\begin{aligned}\mathrm{Vol}(E_4)&=\frac{3}{4}\ r_3\ t_2-\frac{3}{4}\ r_3\ t_4+\frac{3}{4}\ t_2^2-\frac{3}{4}\ t_4^2+\frac{3}{8}\ \sum_{\gamma=1}^{4}t_{1,\gamma}\ t_4-\frac{3}{8}\ \sum_{\gamma=1}^{4}t_{1,\gamma}\ t_2\ ,\\ \mathrm{Vol}(E_{4,1})&=\frac{3}{2}\ r_3\ t_{2,1}-\frac{3}{2}\ r_3\ t_{4,1}+3\ t_{2,1}^2-3\ t_{4,1}^2+\frac{3}{2}\ \sum_{\gamma=1}^{4}t_{1,1\gamma}\ t_{4,1}-\frac{3}{2}\ \sum_{\gamma=1}^{4}t_{1,1\gamma}\ t_{2,1}\ .\end{aligned}$$

The seven (invariant) planes $D_{i\alpha}$ localized at the fix points are given through the relations [4]:

$$\begin{aligned}
&D_1 = \frac{1}{3}\ (R_1 - E_{2,1} - 4\ E_{4,1} - 2\ E_2 - 8\ E_{4,2}\ ) - \frac{1}{3}\ \sum_{\gamma=1}^{4} (\ 3\ E_{3,\gamma} + 2\ E_{1,\gamma} + E_{1,1\gamma}\ )\ , \\
&D_2 = \frac{1}{3}\ (\ R_2 - E_2 - E_4\ ) - \frac{1}{3}\ \sum_{\gamma=1}^{4} E_{1,\gamma}\ ,\ D_{2,1} = \frac{1}{3}\ (\ R_2 - E_{2,1} - E_{4,1}\ ) - \frac{1}{3}\ \sum_{\gamma=1}^{4} E_{1,1\gamma}\ , \\
&D_{3,\gamma} = R_3 - E_{1,1\gamma} - 2\ E_{1,\gamma} - E_{3,\gamma}\ .
\end{aligned} \tag{5.36}$$

In the superpotential (5.24), we have two sets of contributing divisors $\mathcal{D}_i$: On the seven divisors $\mathcal{D}_{D7} = \{D_1, D_{3,\gamma}, E_{2,1}, E_2\}$, a stack of one $O7$–plane and eight $D7$–branes is wrapped. Gaugino condensation takes place in the $SO(8)$ gauge theory. Therefore, we have $a_j = -\frac{2\pi}{6}$ for the set $\mathcal{D}_{D7}$ of divisors contributing in (5.24). On the other hand, divisors in the set $\mathcal{D}_{D3} = \{D_2, D_{2,1}, E_{1,\gamma}, E_{1,1\gamma}, E_{3\gamma}, E_4, E_{4,1}\}$ can be wrapped by Euclidean $D3$–branes. Since all D and E divisors intersect one of the divisors carrying an $O7$–plane in at least one complex dimension (*cf.* Figures 1 and 2), the condition $\chi_{D3} = 1$ for a non–vanishing instanton contribution in the superpotential (5.24) is always met for the set $\mathcal{D}_{D3}$ of divisors. In total we have 23 contributing divisors and the superpotential (5.24) reads:

$$W = W_0(S,U) + \sum_{\mathcal{D}_i \in \mathcal{D}_{D7}} e^{-2\pi\ \frac{\mathrm{Vol}(\mathcal{D}_i)}{6}} + \sum_{\mathcal{D}_i \in \mathcal{D}_{D3}} e^{-2\pi\ \mathrm{Vol}(\mathcal{D}_i)}\ . \tag{5.37}$$

Now we are ready to stabilize all 19 Kähler moduli $r_i, t_{1,\gamma}, t_{1,1\gamma}, t_2, t_{2,1}, t_{3,\gamma}, t_4$ and $t_{4,1}$. The $D3$–brane charge $Q_{3,tot} = -22$ is completely cancelled by the 3–form flux G_3, given in *Eq.* (5.12). In fact, in the previous subsection we have presented critical points for the dilaton and complex structure moduli corresponding to a set of flux solutions, with $N_{flux} = 44$. For a $\widetilde{W}_0 = -0.34864$, corresponding to the critical points of Table 6, we find the following 23 divisor volumes (measured in string units):

$$\begin{array}{rrrl}
\mathrm{Vol}(D_1) = 4.92087\ , & \mathrm{Vol}(D_{2,1}) = 17.1883\ , & \mathrm{Vol}(D_{2,2}) = 17.9329\ , & \\
\mathrm{Vol}(D_{3,\gamma}) = 35.1656\ , & \mathrm{Vol}(E_2) = 3.55689\ , & \mathrm{Vol}(E_4) = 0.710518\ , & \\
\mathrm{Vol}(E_{2,1}) = 4.84171\ , & \mathrm{Vol}(E_{4,1}) = 0.922548\ , & \mathrm{Vol}(E_{3,\gamma}) = 1.01315\ , & \gamma = 1, \ldots, 4\ , \\
\mathrm{Vol}(E_{1,1\gamma}) = 1.06872\ , & \mathrm{Vol}(E_{1,\gamma}) = 0.884484\ , & &
\end{array} \tag{5.38}$$

corresponding to the sizes of the nineteen Kähler moduli:

$$\begin{aligned}
&r_1 = 3.04765 \quad , \quad r_2 = 2.91779 \quad , \quad r_3 = 4.53928\ , \\
&t_{1,\gamma} = 1.52711 \quad , \quad t_{1,1\gamma} = 0.869367 \quad , \quad t_{3,\gamma} = 0.46524 \quad , \quad \gamma = 1, \ldots, 4\ , \\
&t_{2,1} = 0.443261 \quad , \quad t_2 = 0.663503 \quad , \quad t_4 = 0.967525 \quad , \quad t_{4,1} = 0.634432\ .
\end{aligned} \tag{5.39}$$

The divisor volumes give rise to the total volume $\mathrm{Vol}(X_6) = 115.94$. This is large enough, that one–loop (and higher loop) corrections to the Kähler potential are suppressed, with a string–coupling constant $g_{\text{string}} \sim 0.3$ (*cf.* Subsection 5.3). Furthermore, the divisor volumes (5.38) are large enough to suppress higher order instanton effects (*e.g.* from multi–wrapped instantons) in (5.37), since $e^{-2\pi \text{Vol}(E_2)/6} \sim 0.02$ and even smaller for the other divisors.

The six divisors (5.30) or their corresponding cohomology elements give rise to the non–vanishing bulk 2–forms B_2, C_2 (*cf. Eqs.* (3.10) and (3.17)), with the twelve real scalars b^a, c^a. According to *Eq.* (2.5) the latter are combined into the six complex scalars G^a, defined in *Eq.* (2.5). Following Section 3 for stabilizing the moduli b^a we solve the calibration condition (3.13). For this we need the intersection form $\mathcal{K}_{ab}$ and calculate its determinant for the values (5.39). The matrix $\mathcal{K}_{ab}$ is given by:

$$\mathcal{K}_{ab} = \begin{pmatrix} -2\ r_3+8\ t_2-t_4-\sum\limits_\gamma t_{1,\gamma} & \frac{1}{2}\ r_3-t_2+\frac{1}{4}\ \sum\limits_\gamma t_{1,\gamma} & -t_2+\frac{1}{4}\ t_4 & -t_2+\frac{1}{4}\ t_4 & -t_2+\frac{1}{4}\ t_4 & -t_2+\frac{1}{4}\ t_4 \\ 2\ r_3-t_2+2\ \sum\limits_\gamma t_{1,\gamma} & -\frac{1}{2}\ r_3+t_4-\frac{1}{4}\ \sum\limits_\gamma t_{1,\gamma} & -\frac{1}{4}\ t_4+\frac{1}{4}\ t_2 & -\frac{1}{4}\ t_4+\frac{1}{4}\ t_2 & -\frac{1}{4}\ t_4+\frac{1}{4}\ t_2 & -\frac{1}{4}\ t_4+\frac{1}{4}\ t_2 \\ -t_2+\frac{1}{4}\ t_4 & -\frac{1}{4}\ t_4+\frac{1}{4}\ t_2 & \frac{3}{4}\ t_{1,1} & 0 & 0 & 0 \\ -t_2+\frac{1}{4}\ t_4 & -\frac{1}{4}\ t_4+\frac{1}{4}\ t_2 & 0 & \frac{3}{4}\ t_{1,2} & 0 & 0 \\ -t_2+\frac{1}{4}\ t_4 & -\frac{1}{4}\ t_4+\frac{1}{4}\ t_2 & 0 & 0 & \frac{3}{4}\ t_{1,3} & 0 \\ -t_2+\frac{1}{4}\ t_4 & -\frac{1}{4}\ t_4+\frac{1}{4}\ t_2 & 0 & 0 & 0 & \frac{3}{4}\ t_{1,4} \end{pmatrix} . \tag{5.40}$$

For the values (5.39) we find $\det(\mathcal{K}) = -91.25 \neq 0$. Hence, the only non–trivial solution for (3.13) is:

$$b^a = 0\ .$$

Furthermore, in Section 3 a mechanism has been proposed to also stabilize the fields c^a by turning on the 2–form flux ${}^Y f_2$ from the ambient space Y_6 (*cf. Eq.* (3.19) and also the discussion at the end of section 4). To this end, for the $\mathbf{Z}_6$ orbifold we have stabilized all 27 moduli fields.

(*ii*) $\mathbf{Z}_2 \times \mathbf{Z}_4$ – orbifold on the $SU(2)^2 \times SO(5)^2$ lattice :

As our second example, we consider the resolved $\mathbf{Z}_2 \times \mathbf{Z}_4$ orbifold Y_6 on the lattice $SU(2)^2 \times SO(5)^2$. This orbifold has $h_{(1,1)}(Y_6) = 61$ Kähler moduli. We summarize here the relevant data from [4].

The configuration of the fixed point set is displayed in Figure 3 in a schematic way.

There are 16 local $\mathbf{C}^3/\mathbf{Z}_2 \times \mathbf{Z}_4$ patches. The resolution of the $\mathbf{C}^3/\mathbf{Z}_2 \times \mathbf{Z}_4$ singularity is described by the toric diagram in Figure 4. There are four $\mathbf{C}^2/\mathbf{Z}_4$ fixed lines in the z^1 direction from the order four element θ^2. Furthermore, there are $12 + 12 + (10 - 4) = 30$ $\mathbf{C}^2/\mathbf{Z}_2$ fixed lines from the order two elements: From θ^1, $\theta^1(\theta^2)^2$, and $(\theta^2)^2$ in the z^2, z^3, and z^1 direction, respectively. The intersection points of three $\mathbf{Z}_2$ fixed lines are locally described by the resolved $\mathbf{C}^3/\mathbf{Z}_2 \times \mathbf{Z}_2$ patches.



Fig. 3: *Schematic picture of the fixed set configuration of* $\mathbf{Z}_2 \times \mathbf{Z}_4$



Fig. 4: *Toric diagram of two of the resolutions of* $\mathbf{C}^3/\mathbf{Z}_2 \times \mathbf{Z}_4$ *and dual graphs*

The resolution is described by the toric diagram in Figure 5. From these two figures, we can read off the exceptional divisors [4], which together with the inherited divisors R_i form a basis for $H^{(1,1)}(Y_6)$:

$$R_1 \ , \ R_2 \ , \ R_3 \ , \ E_{1,\alpha\gamma'} \ , \ E_{2,\beta\gamma} \ , \ E_{3,\mu} \ , \ E_{4,\beta\gamma} \ , \ E_{5,\alpha\beta\gamma} \ , \ E_{6,\alpha\beta'}, \tag{5.41}$$

with $\alpha = 1, \ldots, 4$, $\beta = 1, 2$, $\beta' = 1, 2, 3$, $\gamma = 1, 2$, $\gamma' = 1, 2, 3$ and $\mu = 1, \ldots, 10$. The divisors $E_{3,\mu}$, $\mu = 1, 2, 4, 5$ will also be denoted by $E_{3,\beta\gamma}$, $\beta, \gamma = 1, 2$. In addition, the orbifold fixed points give rise to the ten divisors [4]:

$$D_{1,\alpha} \quad , \quad D_{2,\beta'} \quad , \quad D_{3,\gamma'} \ . \tag{5.42}$$



Fig. 5: *Toric diagram of the resolution of* $\mathbf{C}^3/\mathbf{Z}_2\times\mathbf{Z}_2$ *and its dual graph*

The topology of these divisors was determined in [4]. The divisors $E_{1,\alpha\gamma}$ and $E_{6,\alpha\beta}$ are blow–ups of $\mathbf{P}^1\times\mathbf{P}^1$ in 5 points, the divisors $E_{1,\alpha 3}$, $E_{3,\mu}$, $\mu=3,6,\ldots,10$, $E_{6,\alpha 3}$, and $D_{1\alpha}$ are blow–ups of $\mathbf{P}^1\times\mathbf{P}^1$ in 4 points, the divisors $E_{3,\mu}$, $\mu=1,2,4,5$ are blow–ups of $\mathbf{P}^1\times\mathbf{P}^1$ in 8 points. The divisors $E_{2,\beta\gamma}$, $E_{4,\beta\gamma}$, $D_{2\beta'}$, and $D_{3\gamma'}$ are $\mathbf{P}^1\times\mathbf{P}^1$, while the divisors $E_{5,\alpha\beta\gamma}$ are $\mathbf{F}_1$, and the R_i are K3 surfaces.

The orientifold projection $\mathcal{O}$ leaves all the divisors (5.41) and (5.42) invariant, hence $h^{(-)}_{(1,1)}(X_6)=0$ (*cf.* also Table 4). We choose an orientifold action σ such that its fixed point set consists of 14 $O7$–planes wrapped on the divisors $D_{1,\alpha}$, $D_{2\beta'}$, $D_{3,\gamma'}$, and $E_{3,\mu}$, $\mu=1,2,4,5$. There are no $O3$–planes. Because of $\chi(D_{1\alpha})=8$, $\chi(D_{2\beta'})=\chi(D_{3,\gamma})=4$, and $\chi(E_{3,\mu})=12$, the total $D3$–brane charge $Q_{3,tot}$ in *Eq.* (5.2) is $Q_{3,tot}=-26$. The Poincare dual 2–forms ω_i of the 61 invariant divisors (5.41) represent a basis for the Kähler form J:

$$J=r_1\ R_1+r_2\ R_2+r_3\ R_3-\sum_{\beta,\gamma=1,2}\left(t_{2,\beta\gamma}\ E_{2,\beta\gamma}+t_{4,\beta\gamma}\ E_{4,\beta\gamma}+\sum_{\alpha=1}^{4}t_{5,\alpha\beta\gamma}\ E_{5,\alpha\beta\gamma}\right)$$
$$-\sum_{\alpha=1}^{4}\left(\sum_{\gamma=1,2,3}t_{1,\alpha\gamma}\ E_{1,\alpha\gamma}+\sum_{\beta=1,2,3}t_{6,\alpha\beta}\ E_{6,\alpha\beta}\right)-\sum_{\mu=1}^{10}t_{3,\mu}\ E_{3,\mu}\ , \tag{5.43}$$

with the 61 Kähler coordinates r_i, $t_{1,\alpha\gamma}$, $t_{2,\beta\gamma}$, $t_{3,\mu}$, $t_{4,\beta\gamma}$, $t_{5,\alpha\beta\gamma}$, $t_{6,\alpha\beta}$. The orientifold action changes the triple intersection numbers $\mathcal{K}_{ijk}$. They become [4]:

$$\begin{aligned}
R_1R_2R_3 &= 2, \qquad R_1E^2_{2\beta\gamma} = -1, \qquad R_1E^2_{3\mu''} = -1,\\
R_1E^2_{4\beta\gamma} &= -1, \qquad R_2E^2_{1\alpha\gamma} = -1, \qquad R_2E^2_{1\alpha 3} = -2,\\
R_3E^2_{6\alpha\beta} &= -1, \qquad R_3E^2_{6\alpha 3} = -2, \qquad E^3_{1\alpha\gamma} = 3/2,\\
E^3_{1\alpha 3} &= 2, \qquad E^2_{1\alpha\gamma'}E_{3\mu''} = -1/2, \qquad E^2_{1\alpha\gamma}E_{6\alpha 3} = -1/2,\\
E^2_{1\alpha 3}E_{6\alpha\beta} &= -1/2, \qquad E^2_{1\alpha 3}E_{6\alpha 3} = -1, \qquad E_{1\alpha\gamma}E^2_{2\beta\gamma} = -1,\\
E_{1\alpha\gamma'}E^2_{3\mu''} &= -1/2, \qquad E_{1\alpha 3}E_{3\mu_1}E_{6\alpha\beta} = 1/2, \qquad E_{1\alpha\gamma}E_{3\mu_2}E_{6\alpha 3} = 1/2,\\
E_{1\alpha 3}E_{3\mu_3}E_{6\alpha 3} &= 1/2, \qquad E_{1\alpha\gamma}E^2_{5\alpha\beta\gamma} = -1, \qquad E_{1\alpha 3}E^2_{6\alpha\beta} = -1/2,\\
E_{1\alpha\gamma}E^2_{6\alpha 3} &= -1/2, \qquad E_{1\alpha 3}E^2_{6\alpha 3} = -1, \qquad E^3_{2\beta\gamma} = 4,\\
E^3_{3\mu''} &= 2, \qquad E^2_{3\mu''}E_{6\alpha\beta'} = -1/2, \qquad E_{3\mu''}E^2_{6\alpha\beta'} = -1/2,\\
E^3_{4\beta\gamma} &= 4, \qquad E^2_{4\beta\gamma}E_{6\alpha\beta} = -1, \qquad E^3_{5\alpha\beta\gamma} = 4,\\
E^2_{5\alpha\beta\gamma}E_{6\alpha\beta} &= -1, \qquad E^3_{6\alpha\beta} = 3/2, \qquad E^3_{6\alpha 3} = 2\ .
\end{aligned} \tag{5.44}$$

The intersection numbers involving the $E_{3,\mu},\ \mu = 1, 2, 4, 5$, which are fixed are

$$\begin{aligned}
R_1E_{3\mu}E_{4\beta\gamma} &= 1, \qquad R_1E_{2\beta\gamma}E_{3\mu} = 1, \qquad E^2_{1\alpha\gamma'}E_{3\mu} = -1,\\
E_{1\alpha\gamma}E_{2\beta\gamma}E_{3\mu} &= 1, \qquad E_{1\alpha\gamma'}E^2_{3\mu} = -2, \qquad E_{1\alpha\gamma}E_{3\mu}E_{5\alpha\beta\gamma} = 1,\\
E^2_{2\beta\gamma}E_{3\mu} &= -4, \qquad E_{2\beta\gamma}E^2_{3\mu} = 4, \qquad E^2_{3\mu}E_{4\beta\gamma} = 4,\\
E^2_{3\mu}E_{6\alpha\beta'} &= -2, \qquad E_{3\mu}E^2_{4\beta\gamma} = -4, \qquad E_{3\mu}E_{4\beta\gamma}E_{6\alpha\beta} = 1,\\
E_{3\mu}E^2_{5\alpha\beta\gamma} &= -2, \qquad E_{3\mu}E_{5\alpha\beta\gamma}E_{6\alpha\beta} = 1, \qquad E_{3\mu}E^2_{6\alpha\beta'} = -1\ .
\end{aligned} \tag{5.45}$$

Furthermore, the orientifold action changes the linear relations [4] between the divisors D_i and R_i:

$$\begin{aligned}
R_1 &= 2\,D_{1,\alpha} + \frac{1}{2}\sum_{\gamma=1}^{3} E_{1,\alpha\gamma} + \frac{1}{2}\sum_{\beta,\gamma=1,2} E_{5,\alpha\beta\gamma} + \frac{1}{2}\sum_{\beta=1}^{3} E_{6,\alpha\beta} \quad , \quad \alpha = 1, \ldots, 4\ ,\\
R_2 &= 4\,D_{2,\beta} + \sum_{\gamma=1,2}\left(\frac{1}{2}\ E_{2,\beta\gamma} + 2\ E_{3,\beta\gamma} + \frac{3}{2}\ E_{4,\beta\gamma}\right) + \frac{1}{2}\sum_{\alpha=1}^{4}\sum_{\gamma=1,2} E_{5,\alpha\beta\gamma}\\
&\quad + \sum_{\alpha=1}^{4} E_{6,\alpha\beta} + E_{3,\mu} \quad , \quad (\beta,\mu) \in \{\ (1,3)\ ,\ (2,6)\ \}\ ,\\
R_2 &= 2\,D_{2,3} + \frac{1}{2}\sum_{\alpha=1}^{4} E_{6,\alpha 3} + \frac{1}{2}\sum_{\mu=7}^{10} E_{3,\mu}\ ,
\end{aligned} \tag{5.46}$$

$$\begin{aligned} R_3 &= 4\, D_{3,\gamma} + \sum_{\alpha=1}^{4} E_{1,\alpha\gamma} + \sum_{\beta=1,2} \left(\frac{3}{2}\ E_{2,\beta\gamma} + 2\ E_{3,\beta\gamma} + \frac{1}{2}\ E_{4,\beta\gamma} \right) + \frac{1}{2} \sum_{\alpha=1}^{4} \sum_{\beta=1,2} E_{5,\alpha\beta\gamma} \\ &\quad + E_{3,\mu} \quad , \quad (\gamma,\mu) \in \{\ (1,7)\ ,\ (2,8)\ \}\ , \\ R_3 &= 2\, D_{3,3} + \frac{1}{2} \sum_{\alpha=1}^{4} E_{1,\alpha 3} + \frac{1}{2} \sum_{\mu=3,6,9,10} E_{3,\mu}\ . \end{aligned}$$

With the intersection numbers (5.44), (5.45) and the relation (5.25), the divisor volumes $\mathrm{Vol}(E)$ and $Vol(D)$ of the 68 divisors (5.41) and (5.42) may be calculated. Since the expressions are rather long we do not display them here.

In the superpotential (5.24), the 68 divisors split into two sets of contributing divisors $\mathcal{D}_i$: On the 14 divisors $\mathcal{D}_{D7} = \{D_{1,\alpha},\ D_{2,\beta},\ D_{3,\gamma},\ E_{3,1},\ E_{3,2},\ E_{3,4},\ E_{3,5}\}$, a stack of one $O7$–plane and eight $D7$–branes is wrapped. Gaugino condensation takes place in the pure $SO(8)$ gauge theory. Therefore we have again $a_j = -\frac{2\pi}{6}$ for the set $\mathcal{D}_{D7}$ of divisors contributing in (5.24). On the other hand, Euclidean $D3$–branes can be wrapped on the divisors in the set $\mathcal{D}_{D3} = \{E_{3,3}, E_{3,6}, E_{3,7}, E_{3,8}, E_{3,9}, E_{3,10}, E_{1,\alpha\gamma}, E_{6,\alpha\beta}, E_{2,\beta\gamma}, E_{4,\beta\gamma}, E_{5,\alpha\beta\gamma}\}$ of the 54 remaining divisors. Since all D and E divisors intersect one of the divisors carrying an $O7$–plane in at least one complex dimension, the condition $\chi_{D3} = 1$ for a non–vanishing instanton contribution in the superpotential (5.24) is always met for the set $\mathcal{D}_{D3}$ of divisors (*cf.* Section 4). In total we obtain for full the superpotential (5.24):

$$W = W_0(S,U) + \sum_{\mathcal{D}_i \in \mathcal{D}_{D7}} e^{-2\pi\ \frac{\mathrm{Vol}(\mathcal{D}_i)}{6}} + \sum_{\mathcal{D}_i \in \mathcal{D}_{D3}} e^{-2\pi\ \mathrm{Vol}(\mathcal{D}_i)}\ . \tag{5.47}$$

Now we are ready to stabilize all 61 Kähler moduli $r_i,\ t_{1,\alpha\gamma},\ t_{2,\beta\gamma},\ t_{3,\mu},\ t_{4,\beta\gamma},\ t_{5,\alpha\beta\gamma},\ t_{6,\alpha\beta}$. The $D3$–brane charge $Q_{3,tot} = -26$ is completely cancelled by the 3–form flux G_3, given in *Eq.* (5.16). In fact, in the previous subsection we have presented critical points for the dilaton and complex structure moduli corresponding to a set of flux solutions, with $N_{flux} = 52$. For a $\widetilde{W_0} = -0.3104$ corresponding to the critical points of Table 8 we find the following 68 divisor volumes (measured in string units):

$$\begin{aligned} \mathrm{Vol}(D_{1,\alpha}) &= 14.00 \quad , \quad \mathrm{Vol}(D_{2,3}) = 5.543 \quad , \quad \mathrm{Vol}(D_{3,3}) = 5.60\ , \\ \mathrm{Vol}(D_{2,\gamma}) &= 10.30 \quad , \quad \mathrm{Vol}(D_{3,\gamma}) = 11.07\ \ , \ \ \gamma = 1,2\ , \\ \mathrm{Vol}(E_{5,\alpha\beta\gamma}) &= 1.30 \quad , \quad \mathrm{Vol}(E_{4,\beta\gamma}) = 1.59\ , \\ \mathrm{Vol}(E_{2,\beta\gamma}) &= 3.30\ \ , \ \ \alpha = 1,\ldots,4\ ,\ \beta,\gamma = 1,2\ , \\ \mathrm{Vol}(E_{1,\alpha\gamma}) &= 9.82 \quad , \quad \mathrm{Vol}(E_{6,\alpha\gamma}) = 14.72\ \ , \ \ \alpha = 1,\ldots,4\ ,\ \gamma = 1,2\ , \\ \mathrm{Vol}(E_{1,\alpha 3}) &= 15.23 \quad , \quad \mathrm{Vol}(E_{6,\alpha 3}) = 21.62\ \ , \ \ \alpha = 1,\ldots,4\ , \\ \mathrm{Vol}(E_{3,\mu}) &= 8.06\ \ , \ \ \mu = 1,2,4,5 \quad , \quad \mathrm{Vol}(E_{3,\mu}) = 27.15\ \ , \ \ \mu = 3,6\ , \\ \mathrm{Vol}(E_{3,\mu}) &= 19.26\ \ , \ \ \mu = 7,8 \quad , \quad \mathrm{Vol}(E_{3,\mu}) = 27.00\ \ , \ \ \mu = 9,10\ . \end{aligned} \tag{5.48}$$

corresponding to the sizes of the 61 Kähler moduli:

$$\begin{aligned} r_1 &= 7.826 \quad , \quad r_2 = 5.410 \quad , \quad r_3 = 4.593 \ , \\ t_{5,\alpha\beta\gamma} &= 0.770 \quad , \quad t_{4,\beta\gamma} = 0.107 \ , \\ t_{2,\beta\gamma} &= 0.239 \quad , \quad \alpha = 1, \ldots, 4 \ , \ \beta, \gamma = 1, 2 \ , \\ t_{1,\alpha\gamma} &= 0.858 \quad , \quad t_{6,\alpha\gamma} = 1.860 \quad , \quad \alpha = 1, \ldots, 4 \ , \ \gamma = 1, 2 \ , \\ t_{1,\alpha 3} &= 0.686 \quad , \quad t_{6,\alpha 3} = 1.320 \quad , \quad \alpha = 1, \ldots, 4 \ , \\ t_{3,\mu} &= 0.037 \quad , \quad \mu = 1, 2, 4, 5 \quad , \quad t_{3,\mu} = 0.569 \quad , \quad \mu = 3, 6 \ , \\ t_{3,\mu} &= 0.302 \quad , \quad \mu = 7, 8 \quad , \quad t_{3,\mu} = 0.413 \quad , \quad \mu = 9, 10 \ . \end{aligned} \tag{5.49}$$

The divisor volumes give rise to the total volume $\mathrm{Vol}(X_6) = 229.22$. This is large enough, that one–loop corrections to the Kähler potential are suppressed. Furthermore, the divisor volumes (5.48) are large enough to suppress higher order instanton effects in (5.47), since *e.g.* $e^{-2\pi \mathrm{Vol}(E_{5,\alpha\beta\gamma})} \sim 0.0002$ and even smaller for the other divisors. To this end, for the $\mathbf{Z}_2 \times \mathbf{Z}_4$ orbifold we have stabilized all 63 moduli fields.

5.5. Orbifolds without complex structure moduli

It has already been discussed in [24,3] and thoroughly in Subsection 2.4. that type IIB orientifolds of toroidal $\mathbf{Z}_N$ and $\mathbf{Z}_N \times \mathbf{Z}_M$ orbifolds without complex structure are not suitable candidates for a KKLT scenario. This investigation has been limited to the singular orbifold case. According to Table 3, this concerns the following orbifolds: $\mathbf{Z}_3, \mathbf{Z}_7, \mathbf{Z}_3 \times \mathbf{Z}_3$, $\mathbf{Z}_4 \times \mathbf{Z}_4$, $\mathbf{Z}_6 \times \mathbf{Z}_6$, $\mathbf{Z}_2 \times \mathbf{Z}_{6'}$ and $\mathbf{Z}_{8-I}$ on the $SU(4)^2$ lattice. In this subsection we shall extend this discussion to resolved orbifolds. We want to apply our stability criteria, derived in Subsection 2.3, to check whether those orbifolds with the generic superpotential (2.23) may be suitable for a stable uplift.

We consider an orientifold compactification on a resolved orbifold X_6 with $n := h^{(+)}_{(1,1)}(X_6)$ Kähler moduli (and $h^{(-)}_{(2,1)}(X_6) = 0$). There are n_1 untwisted moduli r^i and n_2 blow–up moduli t^α, with $n = n_1 + n_2$. We may introduce the Kähler form $\mathcal{J}$:

$$\mathcal{J} = \sum_{i=1}^{n_1} r^i \ u_i - \sum_{\alpha=1}^{n_2} t^\alpha \ v_\alpha \ , \tag{5.50}$$

with the Kähler coordinates $r^i > 0$ and $t^\alpha > 0$. The volume of X_6 is given by

$$\mathrm{Vol}(X_6) = \frac{1}{6} \int_{X_6} \mathcal{J} \wedge \mathcal{J} \wedge \mathcal{J} \ .$$

We consider the set of n divisors $\widetilde{\mathcal{D}}_i \in \left\{\bigcup_{i=1}^{n_1} R_i \ , \ \bigcup_{\alpha=1}^{n_2} \widetilde{E}_\alpha\right\}$, which are Poincaré dual to the classes $\widetilde{\omega}_i \in \{\bigcup_{i=1}^{n_1} u_i \ , \ \bigcup_{\alpha=1}^{n_2} v_\alpha\}$. Their volumes are given as usual by:

$$\begin{aligned} V_{r^i} &= \frac{3}{2}\ \frac{\partial}{\partial r^i}\ \mathrm{Vol}(X_6) \quad , \quad i = 1, \ldots, n_1 \\ V_{t^\alpha} &= \frac{3}{2}\ \frac{\partial}{\partial t^\alpha}\ \mathrm{Vol}(X_6) \quad , \quad \alpha = 1, \ldots, n_2 \ . \end{aligned} \tag{5.51}$$

Following the discussion in Subsection 2.3, the divisors $\widetilde{\mathcal{D}}_i$ do not yet necessarily represent contributing divisors $\mathcal{D}_i$ in (2.23). However, for resolved orbifolds X_6, the latter are generically related to the n original divisors $\widetilde{\mathcal{D}}_i$ through linear combinations of the form

$$\begin{aligned} D_i &= \lambda_i\ R_i + \sum_{\alpha=1}^{n_2} \mu_{i\alpha}\ \widetilde{E}_\alpha \quad , \quad i = 1, \ldots, n_1 \ , \\ E_\alpha &= -\rho_\alpha \widetilde{E}_\alpha \quad , \quad \alpha = 1, \ldots, n_2 \ , \end{aligned} \tag{5.52}$$

with some positive rational numbers $\lambda_i, \mu_{i\alpha}, \rho_\alpha \in \mathbf{Q}^+$. Following *Eq.* (2.52), the invertible $n \times n$–matrix N assumes the form

$$N = \begin{pmatrix} \lambda & \mu \\ 0 & -\rho \end{pmatrix} \ , \tag{5.53}$$

with the submatrices

$$\lambda = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_{n_1} \end{pmatrix} \quad , \quad \mu = \begin{pmatrix} \mu_{11} & \ldots & \mu_{1n_2} \\ \vdots & & \vdots \\ \mu_{n_1 1} & \ldots & \mu_{n_1 n_2} \end{pmatrix} \quad , \quad \rho = \begin{pmatrix} \rho_1 & & \\ & \ddots & \\ & & \rho_{n_2} \end{pmatrix} \ .$$

According to *Eq.* (2.53), the n new Kähler coordinates corresponding to the contributing divisors (5.52) are related to the original orbifold coordinates r^i, t^α by the matrix $(N^t)^{-1}$, *i.e.* :

$$\begin{pmatrix} \lambda^{-1} & 0 \\ \rho^{-1}\ \mu^t\ \lambda^{-1} & -\rho^{-1} \end{pmatrix} \begin{pmatrix} r^1 \\ \vdots \\ r^{n_1} \\ t^1 \\ \vdots \\ t^{n_2} \end{pmatrix} \ . \tag{5.54}$$

Note, that the matrices λ^{-1} and $\rho^{-1}\ \mu^t\ \lambda^{-1}$ have only positive entries and the negative $n_2 \times n_2$–matrix $-\rho^{-1}$ only acts on the blow–up moduli t^α, which generically are very small. Hence all the n new Kähler coordinates, corresponding to the divisors (5.52), are positive, *i.e.* the condition (2.45) is fulfilled.

For concreteness, let us discuss an example. The (symmetrically) resolved $\mathbf{Z}_2 \times \mathbf{Z}_2$ orientifold has 3 untwisted Kähler moduli corresponding to the three radii r^i of the six–torus $T^6 = (T^2)^3$ and 48 twisted Kähler moduli t^i describing the blow–up modes or exceptional divisors of the orbifold singularities. In the limit of $r_i = r$ and $t_i = t$, the volume of this orientifold becomes [16]

$$Vol(X_6) = r^3 - 24\ r\ t^2 + 48\ t^3\ . \tag{5.55}$$

From $\mathcal{K} = 6Vol(X_6)$ the intersection numbers $\widetilde{\mathcal{K}}_{ijk}$ may be read off

$$\widetilde{\mathcal{K}}_{rrr} = 6 \quad , \quad \widetilde{\mathcal{K}}_{rtt} = -48 \quad , \quad \widetilde{\mathcal{K}}_{ttt} = 288\ , \tag{5.56}$$

with the intersection matrix

$$\widetilde{\mathcal{K}}_{ij} = 6\ \begin{pmatrix} r & -8\ t \\ -8\ t & 48\ t - 8\ r \end{pmatrix}\ . \tag{5.57}$$

With the relation $V_{\widetilde{\mathcal{D}}_i} = \frac{3}{4}\widetilde{\mathcal{K}}_{ijk} t^j t^k$ we calculate the divisor volumes $V_{\widetilde{\mathcal{D}}_i}$ of the divisors $\widetilde{\mathcal{D}}_i$ corresponding to the Kähler moduli $\widetilde{t}^1 = r$ and $\widetilde{t}^2 = t$:

$$V_{\widetilde{\mathcal{D}}_1} = \frac{9}{2}\ (r^2 - 8t^2) \quad , \quad V_{\widetilde{\mathcal{D}}_2} = 72\ t\ (3t - r)\ . \tag{5.58}$$

The divisors $\mathcal{D}_i$, which contribute to the superpotential (2.23), are constructed from the divisors $\widetilde{\mathcal{D}}_i$ through a linear combination (2.52). The volume of the right divisors $\mathcal{D}_i$, representing also the real parts of the holomorphic Kähler moduli T^1, T^2 become [16]:

$$\begin{aligned} \mathrm{Re}(T^1) &:= V_{\mathcal{D}_1} = \frac{1}{3}\ V_{\widetilde{\mathcal{D}}_1} + \frac{8}{48}\ V_{\widetilde{\mathcal{D}}_2} = \frac{3}{2}\ (r - 4t)^2\ , \\ \mathrm{Re}(T^2) &:= V_{\mathcal{D}_2} = -\frac{1}{48}\ V_{\widetilde{\mathcal{D}}_2} = \frac{3}{2}\ t\ (r - 3t)\ . \end{aligned} \tag{5.59}$$

In terms of the latter, the Kähler potential $K_{KM} = -2\ln Vol(X_6)$ for the Kähler moduli T^1, T^2 becomes:

$$\begin{aligned} K = {} & 2\ln(T^2 + \overline{T}^2) - \ln\Big\{(T^1 + \overline{T}^1) + 2\ (T^2 + \overline{T}^2) \\ & -(T^1 + \overline{T}^1)^{1/2}\ [T^1 + \overline{T}^1 + 4\ (T^2 + \overline{T}^2)]^{1/2}\Big\} \\ & - 2\ln\Big\{(T^1 + \overline{T}^1)^2 + 16\ (T^1 + \overline{T}^1)\ (T^2 + \overline{T}^2) + 32\ (T^2 + \overline{T}^2)^2 \\ & + (T^1 + \overline{T}^1)^{3/2}\ [(T^1 + \overline{T}^1) + 4\ (T^2 + \overline{T}^2)]^{1/2} \\ & +8\ (T^2 + \overline{T}^2)\ (T^1 + \overline{T}^1)^{1/2}\ [(T^1 + \overline{T}^1) + 4\ (T^2 + \overline{T}^2)]^{1/2}\Big\}\ . \end{aligned} \tag{5.60}$$

The matrix N, defined in (2.52), is:

$$N = \begin{pmatrix} \frac{1}{3} & \frac{1}{6} \\ 0 & -\frac{1}{48} \end{pmatrix} \ . \tag{5.61}$$

According to (2.53) the Kähler coordinates t^i related to the contributing divisors $\mathcal{D}_i$ are[11]:

$$t^1 = 3\ r > 0 \quad , \quad t^2 = 24\ (r - 2t) > 0\ . \tag{5.63}$$

The positivity properties follow from the condition, that r, t lie in the Kähler cone $r > 4t$ [16]. Expressed in terms of the coordinates t^i, the volume $Vol(X_6)$ reads:

$$\mathcal{K} = \frac{2}{9}\ (t^1)^3 - \frac{1}{6}\ (t^1)^2\ t^2 + \frac{1}{24}\ t^1\ (t^2)^2 - \frac{1}{384}\ (t^2)^3\ . \tag{5.64}$$

Finally, the intersection form $\mathcal{K}_{ij} = N \widetilde{\mathcal{K}}_{ij} N^t$ becomes:

$$\mathcal{K}_{ij} = \begin{pmatrix} \frac{2}{9} t^1 - \frac{1}{18}\ t^2 & -\frac{1}{18} t^1 + \frac{1}{72}\ t^2 \\ -\frac{1}{18} t^1 + \frac{1}{72}\ t^2 & \frac{1}{72}\ t^1 - \frac{1}{384} t^2 \end{pmatrix} \ . \tag{5.65}$$

Now we may apply our condition (2.45) to verify whether the ansatz (2.23) with the contributing divisors $\mathcal{D}_i$ may lead to a stable uplift. The criterion (2.45) is met, since $t^1 > 0$ and $t^2 > 0$. Hence, no uplift is possible with the ansatz (2.23). This means that a stable minimum may be only found in the $\mathbf{Z}_2 \times \mathbf{Z}_2$ orientifold if one also includes the minimization w.r.t. the complex structure moduli [16].

Similarly, the condition (2.45) may be verified for the orbifolds $\mathbf{Z}_3, \mathbf{Z}_7, \mathbf{Z}_3 \times \mathbf{Z}_3$, $\mathbf{Z}_4 \times \mathbf{Z}_4$, $\mathbf{Z}_6 \times \mathbf{Z}_6$, $\mathbf{Z}_2 \times \mathbf{Z}_{6'}$, and $\mathbf{Z}_8 - I$ with $SU(4)^2$ lattice to conclude, that these manifolds are not appropriate for a KKLT scenario.

6. Concluding remarks

In this article we have discussed moduli stabilization in type IIB CY orientifold compactifications within the approach of the KKLT scenario. In Section 2 we have derived general conditions for a stable uplift following from the F–flatness conditions (2.17). We have investigated the stability criteria of type IIB CY orientifold compactifications without

---

[11] According to (2.44) these coordinates may be read off from the first derivatives of the Kähler potential K:

$$t^1 = -3V\ K_{T^1} = 3\ r \quad , \quad t^2 = -3V\ K_{T^2} = 24\ (r - 2t)\ . \tag{5.62}$$

complex structure moduli, *i.e.* $h^{(-)}_{(2,1)}(X_6) = 0$. For this class of compactifications, we have derived general criteria on the Kähler potential, *cf. Eqs.* (2.36), (2.40), (2.45) or (2.46). If one of these criteria holds, the axionic mass matrix is not positive definite and no stable uplift is possible. In particular, the latter conditions are met in the type IIB orientifolds of toroidal orbifold compactifications without complex structure moduli: $\mathbf{Z}_3, \mathbf{Z}_7, \mathbf{Z}_3 \times \mathbf{Z}_3$, $\mathbf{Z}_4 \times \mathbf{Z}_4$, $\mathbf{Z}_6 \times \mathbf{Z}_6$, $\mathbf{Z}_2 \times \mathbf{Z}_{6'}$, and $\mathbf{Z}_8 - I$ with $SU(4)^2$ lattice. This is true both at the orbifold point and after the resolution. Hence, no stable uplift is possible for those orbifolds, *cf.* Subsection 5.5. In fact, all the existing type IIB orientifolds without complex structure moduli fulfill one of the conditions *Eqs.* (2.36), (2.40), (2.45) or (2.46) and therefore compactifications without complex structure moduli generically seem not to allow a stable uplift. Furthermore, in Section 3 we have presented a mechanism to stabilize Kähler moduli from the cohomology $H^{(2)}_{-}(X_6)$ by analyzing the calibration conditions of 4–cycles. Equipped with these results we accomplished to fix all moduli in some examples of resolved orbifolds: $\mathbf{Z}_{6-II}$ on the root lattice of $SU(2) \times SU(6)$ and $\mathbf{Z}_2 \times \mathbf{Z}_4$ on $SU(2)^2 \times SO(5)^2$: We have stabilized for the $\mathbf{Z}_{6-II}$–orbifold all its 27 moduli fields, *cf.* Table 4 and (5.39) and for the $\mathbf{Z}_2 \times \mathbf{Z}_4$ orbifold all its 63 moduli fields, *cf.* Table 6 and (5.49).

In most of the existing literature on flux compactifications, one works in the lowest α'–expansion at string tree–level, *i.e.* in the supergravity approximation. Since at this order in α' and g_{string} the theory has a no–scale structure, which does not fix all Kähler moduli, one adds some effects which eventually allow the stabilization of all moduli fields. As proposed by KKLT [1], one promising possibility is to consider the racetrack superpotential (5.24), which we also have used throughout this article. The critical points found in Subsection 5.4 are inert against corrections in α' and g_{string}, since the coupling constant g_{string} is small and the volume V of the compactification manifold is large. However, to discuss also other choices of minima one must go beyond this approximation and include corrections in both α' and g_{string}. In particular, there are both perturbative corrections to the Kähler potential K to all orders in g_{string} and world–sheet as well as space–time instanton corrections to the Kähler potential K in N=1 CY orientifolds. It is certainly very important to gain control over these corrections.

The coefficient $\gamma_j(S,U)$ in the non–perturbative superpotential (5.24) accounts for gaugino condensation or $D3$–instanton effects. Generically, this weight factor depends both on the dilaton S and the complex structure moduli U^λ. Due to the non–renormalization of the gauge kinetic function beyond one–loop, for gaugino condensation this dependence is fairly well under control perturbatively, *cf. Eq.* (2.15). On the other hand, for $D3$–instantons the factor $\gamma_j(S,U)$ represents the one–loop determinant of the instanton solution. The latter is hard to compute directly, except (in)directly in F– or M–theory [9] or through some duality arguments [54,55]. In Section 4 we have presented general results (*cf.* Table 2), under which conditions this coefficient γ is non–vanishing in type IIB

CY orientifolds. However, it is certainly important to directly calculate the dilaton and complex structure modulus dependence of γ by means of an instanton calculation.

We have obtained a fairly complete picture of the critical points of type IIB orientifolds of resolved orbifolds. Indeed, as Tables 6 and 8 show, for one orbifold there is a huge number of vacua with the same physical quantities thus giving rise to a landscape of supersymmetric vacuum solutions in the flux space. Throughout this article, the fixing of open string moduli is not addressed. This is legitimate as we only discuss $D3$–branes wrapping internal 4–cycles and no space–time filling $D3$–branes. The complete tadpole (5.2) originating from the RR 4–form is cancelled by curvature and flux. More general setups would also allow space–time filling $D3$–branes and $D7$–branes away from the orientifold planes. It has been shown in [49,56], that even an ISD 3–form flux implies stabilization of the $D7$–brane positions and soft–masses for corresponding the open string moduli. Certainly, a thorough discussion of the stabilization of open string moduli would enrich the present picture of the string landscape [57].

Acknowledgments

We are indebted to Hans Jockers and Peter Mayr for many valuable discussions. Moreover we wish to thank Thomas Grimm, Renata Kallosh, and Alexander Schmidt for useful discussions.

This work is supported in part by the Deutsche Forschungsgemeinschaft as well as by the EU-RTN network *Constituents, Fundamental Forces and Symmetries of the Universe* (MRTN-CT-2004-005104) and is a collaboration between the contractors Ludwig–Maximilians University at Munich and Universita di Torino to which the Universita del Piemonte Orientale, Alessandria, is associated.

S.R. and W.S. thank the university of Munich for hospitality. The work of E. S. is supported by the Marie Curie Grant MERG–CT–2004–006374. E.S. thanks the Erwin Schroedinger Institute in Vienna, the Simons Workshop in Mathematics and Physics at Stony Brook, the Arnold Sommerfeld Center for Theoretical Physics and the Ludwig–Maximilians University at Munich for hospitality.

Appendix A. Complex structures of $\mathbf{Z}_N-$ and $\mathbf{Z}_N \times \mathbf{Z}_M$–orbifolds

In this appendix we introduce a complex basis and complex structures for the orbifolds $\mathbf{Z}_{6-II}$ on $SU(2) \times SU(6)$, $\mathbf{Z}_2 \times \mathbf{Z}_4$ on $SU(2)^2 \times SO(5)^2$ and $\mathbf{Z}_4$ on $SU(4)^2$.

(i) $\mathbf{Z}_{6-II}$ – orbifold with $SU(2) \times SU(6)$ lattice

We introduce the complex coordinates and complex structures [3]:

$$\begin{aligned} dz^1 &= dx^1 + e^{2\pi i/6} dx^2 + e^{2\pi i/3} dx^3 - dx^4 + e^{-2\pi i/3} dx^5 \ , \\ dz^2 &= dx^1 + e^{2\pi i/3} dx^2 + e^{-2\pi i/3} dx^3 + dx^4 + e^{2\pi i/3} dx^5 \ , \\ dz^3 &= \frac{1}{\sqrt{12}} \left[\ \frac{1}{3} \ (dx^1 - dx^2 + dx^3 - dx^4 + dx^5) - iU \ dx^6 \ \right] \ . \end{aligned} \tag{A.1}$$

$$\begin{aligned} \omega_{A0} &= dz^1 \wedge dz^2 \wedge dz^3 \ , \\ \omega_{B0} &= d\bar{z}^1 \wedge d\bar{z}^2 \wedge d\bar{z}^3 \ , \\ \omega_{A3} &= dz^1 \wedge dz^2 \wedge d\bar{z}^3 \ , \\ \omega_{B3} &= d\bar{z}^1 \wedge d\bar{z}^2 \wedge dz^3 \ . \end{aligned} \tag{A.2}$$

$$\begin{aligned} \int \omega_{A0} \wedge \omega_{B0} &= -i(U + \bar{U}) \ , \\ \int \omega_{A3} \wedge \omega_{B3} &= -i(U + \bar{U}) \ . \end{aligned} \tag{A.3}$$

(ii) $\mathbf{Z}_2 \times \mathbf{Z}_4$ – orbifold with $SU(2)^2 \times SO(5)^2$ lattice :

We introduce the complex coordinates and complex structures:

$$\begin{aligned} dz^1 &= dx^1 + U \ dx^2 \ , \\ dz^2 &= dx^3 - \frac{1}{2} \ (1 - i) \ dx^4 \ , \\ dz^3 &= dx^5 - \frac{1}{2} \ (1 + i) \ dx^6 \ . \end{aligned} \tag{A.4}$$

(iii) $\mathbf{Z}_4$ – orbifold with $SU(4)^2$ lattice

We introduce the complex coordinates and complex structures:

$$\begin{aligned} dz^1 &= dx^1 - dx^3 + i \ dx^2 \ , \\ dz^2 &= dx^4 - dx^6 + i \ dx^5 \ , \\ dz^3 &= dx^1 - dx^2 + dx^3 + U \ (dx^4 - dx^5 + dx^6) \ . \end{aligned} \tag{A.5}$$

In the complex basis we have[12] [58]:

$$\begin{aligned} n^t g n = & \, 2 \ \mathrm{Re}T^1 \ |N^1|^2 + 2 \ \mathrm{Re}T^2 \ |N^2|^2 + 2 \ \mathrm{Re}T^3 \ |N^3|^2 \\ & + 2 \ (\mathrm{Re}T^{12} + i \ \mathrm{Re}T^{21}) \ N^1 \overline{N}^2 + 2 \ (\mathrm{Re}T^{12} - i \ \mathrm{Re}T^{21}) \ \overline{N}^1 \ N^2 \ , \end{aligned} \tag{A.6}$$

with the complex (normalized) lattice vectors

$$\begin{aligned} N^1 &= \frac{1}{\sqrt{2}} \ (n^1 - n^3 + i \ n^2) \ , \\ N^2 &= \frac{1}{\sqrt{2}} \ (n^4 - n^6 + i \ n^5) \ , \\ N^3 &= \frac{1}{\sqrt{2U_2}} \ \frac{1}{2} \ [\ n^1 - n^2 + n^3 + U \ (n^4 - n^5 + n^6) \ ] \end{aligned} \tag{A.7}$$

following from (A.5). The definition of the complex numbers N^i represents an unimodular transformation, *i.e.* $detY = 1$, on the integers n^i. Hence it preserves the volume of the integer space n^i when moving to the complex space N^i. Note the additional factor of 1/2 for N^3. The Kähler moduli follow

$$\begin{aligned} T^1 &= R^2 + x + i \ \alpha \ , \\ T^2 &= S^2 + y + i \ \beta \ , \\ T^3 &= 2 \ \sqrt{4xy - (v+z)^2} + 2 \ i \ (\delta + \epsilon) \ , \\ T^{12} &= u + \frac{1}{2} \ (v+z) + \frac{i}{2} \ (\delta - \epsilon) \ , \\ T^{21} &= \frac{1}{2} \ (v - z) - i \ \gamma - \frac{i}{2} \ (\delta + \epsilon) \ , \end{aligned} \tag{A.8}$$

with

$$(\det g)^{1/2} = \frac{1}{8} \ [(T^1 + \overline{T}^1) \ (T^2 + \overline{T}^2) - (T^{12} + \overline{T}^{12})^2 - (T^{21} + \overline{T}^{21})^2 \ ] \ (T^3 + \overline{T}^3) \ .$$

Furthermore, the complex structure [58]:

$$U = \frac{1}{2 \ x} \ [\ v + z - i \ \sqrt{4xy - (v+z)^2} \ ] \ . \tag{A.9}$$

---

[12] Note that for a 2–torus we have $n^t g n = 2 \ \mathrm{Re}T \ |N|^2$, with $N = \frac{1}{\sqrt{2U_2}} \ (n^1 + U \ n^2)$. Sometimes one also introduces $\widetilde{N} = n^1 + U \ n^2$ such that $n^t g n = 2 \ \mathrm{Re}T \ \left(\frac{1}{\sqrt{2U_2}}\right)^2 \ |\widetilde{N}|^2$. Essentially, the two coordinates N and $\widetilde{N}$ only affect the normalization of the volume (U_2 or $1/2$) of the complex torus with the corners $(0, 1, U, 1 + U)$ or $\frac{1}{\sqrt{2U_2}} \ (0, 1, U, 1 + U)$, respectively. At any rate the coordinate N represents an unimodular transformation from the integer space n^i to the complex space, *i.e.* it preserves the volume of the integer lattice and there is an one to one map from n^i to N.

Following [3] we may introduce new Kähler moduli:

$$\begin{aligned}
\widetilde{T}^{12} &= T^{12}+i\,T^{21} = u+\frac{1}{2}(1+i)\ v+\frac{1}{2}(1-i)\ z+\gamma+\frac{1}{2}(1+i)\ \delta+\frac{1}{2}(1-i)\epsilon\ ,\\
\widetilde{T}^{21} &= T^{12}-i\,T^{21} = u+\frac{1}{2}(1-i)\ v+\frac{1}{2}(1+i)\ z-\gamma-\frac{1}{2}(1-i)\ \delta-\frac{1}{2}(1+i)\epsilon
\end{aligned} \tag{A.10}$$

such that

$$(\mathrm{det}g)^{1/2}=\frac{1}{8}\ [(T^1+\overline{T}^1)\ (T^2+\overline{T}^2)-(\widetilde{T}^{12}+\overline{\widetilde{T}^{21}})\ (\widetilde{T}^{21}+\overline{\widetilde{T}^{12}})\ ]\ (T^3+\overline{T}^3)\ .$$

Clearly any transformation $n^i \longrightarrow n^i+1$ is a symmetry of the T^6 reflecting the periodicity $x^i \longrightarrow x^i+1$ of the lattice (in the lattice basis). It is not a priori guaranteed, that our three complex tori, introduced through the coordinates (A.7) share this periodicity. The third torus lies within the full six–dimensional lattice and has been defined such, that the integers n^i are mapped one to one onto the complex numbers N^i. By introducing new lattice vectors $\widetilde{n}^i$ in (A.7) we may disentangle the six windings n^i such that our three complex tori become independent. We introduce:

$$\begin{aligned}
&\widetilde{n}^1=n^1-n^3 \quad ,\quad \widetilde{n}^2=n^2\ ,\\
&\widetilde{n}^4=n^4-n^6 \quad ,\quad \widetilde{n}^5=n^5\ ,\\
&\widetilde{n}^3=\frac{1}{2}\ (n^1+n^3-n^2) \quad ,\quad \widetilde{n}^6=\frac{1}{2}\ (n^4+n^6-n^5)\ .
\end{aligned} \tag{A.11}$$

Since $\mathrm{det}Y=1$, with $\widetilde{n}^i=Y^i_j\ n^j$, the map (A.11) represents an one to one transformation from the integers n^i to the numbers $\widetilde{n}^i$. The latter however may become half–integer. This way (A.7) becomes:

$$\begin{aligned}
\widetilde{N}^1&=\frac{1}{\sqrt{2}}\ (\widetilde{n}^1+i\ \widetilde{n}^2)\quad ,\quad \widetilde{n}^1,\widetilde{n}^2\in\mathbf{Z}\ ,\\
\widetilde{N}^2&=\frac{1}{\sqrt{2}}\ (\widetilde{n}^4+i\ \widetilde{n}^5)\quad ,\quad \widetilde{n}^4,\widetilde{n}^5\in\mathbf{Z}\ ,\\
\widetilde{N}^3&=\frac{1}{\sqrt{2U_2}}\ (\widetilde{n}^3+U\ \widetilde{n}^6)\quad ,\quad \widetilde{n}^3,\widetilde{n}^6\in\frac{1}{2}\mathbf{Z}\ .
\end{aligned} \tag{A.12}$$

Now, the complex coordinates $\widetilde{N}^i$ assume the expected form of three independent two–tori (*cf.* the above footnote). Note, that the property $\widetilde{n}^3,\widetilde{n}^6\in\frac{1}{2}\mathbf{Z}$ means, that some lattice points n^i are mapped into the third torus rather than onto its corners. E.g. the point $n^6=1$ becomes $\widetilde{n}^6=\frac{1}{2}$. Nevertheless, the third torus with the complex structure U is spanned by the points $(\widetilde{n}^3,\widetilde{n}^6)=(0,0),\ (0,1),\ (0,1)$ and $(1,1)$, with $\widetilde{n}^1,\widetilde{n}^2,\widetilde{n}^4,\widetilde{n}^5=0$. This corresponds to the original lattice points $(n^1,n^2,n^3,n^4,n^5,n^6)=(1,0,1,1,0,1)$. This way, the third torus is decoupled from the other two.

$$\begin{aligned}
dz^1&=dx^1-dx^3+i\ dx^2\ ,\\
dz^2&=dx^4-dx^6+i\ dx^5\ ,\\
dz^3&=\frac{1}{4}\left[\ dx^1-dx^2+dx^3-i\ U\ \left(dx^4-dx^5+dx^6\right)\ \right]\ .
\end{aligned} \tag{A.13}$$

References

[1] S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, "De Sitter vacua in string theory," Phys. Rev. D **68**, 046005 (2003) [arXiv:hep-th/0301240].

[2] C.P. Burgess, R. Kallosh and F. Quevedo, "de Sitter string vacua from supersymmetric D-terms," JHEP **0310**, 056 (2003) [arXiv:hep-th/0309187].

[3] D. Lüst, S. Reffert, W. Schulgin and S. Stieberger, "Moduli stabilization in type IIB orientifolds. I," arXiv:hep-th/0506090.

[4] D. Lüst, S. Reffert, E. Scheidegger, S. Stieberger "Resolved Toroidal Orbifolds and their Orientifolds," arXiv:hep-th/0609014.

[5] S. Reffert and E. Scheidegger, "Moduli stabilization in toroidal type IIB orientifolds," Fortsch. Phys. **54**, 462 (2006) [arXiv:hep-th/0512227].

[6] F. Gmeiner, R. Blumenhagen, G. Honecker, D. Lüst and T. Weigand, "One in a billion: MSSM-like D-brane statistics," JHEP **0601**, 004 (2006) [arXiv:hep-th/0510170].

[7] S. B. Giddings, S. Kachru and J. Polchinski, "Hierarchies from fluxes in string compactifications," Phys. Rev. D **66**, 106006 (2002) [arXiv:hep-th/0105097].

[8] S. Gukov, C. Vafa and E. Witten, "CFT's from Calabi-Yau four-folds," Nucl. Phys. B **584**, 69 (2000) [Erratum-ibid. B **608**, 477 (2001)] [arXiv:hep-th/9906070];
T.R. Taylor and C. Vafa, "RR flux on Calabi-Yau and partial supersymmetry breaking," Phys. Lett. B **474**, 130 (2000) [arXiv:hep-th/9912152];
P. Mayr, "On supersymmetry breaking in string theory and its realization in brane worlds," Nucl. Phys. B **593**, 99 (2001) [arXiv:hep-th/0003198].

[9] E. Witten, "Non-Perturbative Superpotentials In String Theory," Nucl. Phys. B **474**, 343 (1996) [arXiv:hep-th/9604030].

[10] S. Ferrara, L. Girardello and H.P. Nilles, "Breakdown Of Local Supersymmetry Through Gauge Fermion Condensates," Phys. Lett. B **125**, 457 (1983);
M. Dine, R. Rohm, N. Seiberg and E. Witten, "Gluino Condensation In Superstring Models," Phys. Lett. B **156**, 55 (1985).

[11] P. Breitenlohner and D.Z. Freedman, "Stability In Gauged Extended Supergravity," Annals Phys. **144**, 249 (1982).

[12] B. Acharya, M. Aganagic, K. Hori and C. Vafa, "Orientifolds, mirror symmetry and superpotentials," arXiv:hep-th/0202208.

[13] I. Brunner and K. Hori, "Orientifolds and mirror symmetry," JHEP **0411**, 005 (2004) [arXiv:hep-th/0303135].

[14] T.W. Grimm and J. Louis, "The effective action of N = 1 Calabi-Yau orientifolds," Nucl. Phys. B **699**, 387 (2004) [arXiv:hep-th/0403067];
T.W. Grimm, "The effective action of type II Calabi-Yau orientifolds," Fortsch. Phys. **53**, 1179 (2005) [arXiv:hep-th/0507153]

[15] F. Denef, M.R. Douglas and B. Florea, “Building a better racetrack,” JHEP **0406**, 034 (2004) [arXiv:hep-th/0404257].

[16] F. Denef, M.R. Douglas, B. Florea, A. Grassi and S. Kachru, “Fixing All Moduli in a Simple F-Theory Compactification,” arXiv:hep-th/0503124.

[17] L. Görlich, S. Kachru, P.K. Tripathy and S.P. Trivedi, “Gaugino condensation and nonperturbative superpotentials in flux compactifications,” JHEP **0412**, 074 (2004) [arXiv:hep-th/0407130].

[18] D. Lüst, P. Mayr, S. Reffert and S. Stieberger, “F-theory flux, destabilization of orientifolds and soft terms on D7-branes,” Nucl. Phys. B **732**, 243 (2006) [arXiv:hep-th/0501139].

[19] D. Lüst, S. Reffert and S. Stieberger, “MSSM with soft SUSY breaking terms from D7-branes with fluxes,” Nucl. Phys. B **727**, 264 (2005) [arXiv:hep-th/0410074].

[20] K. Choi, A. Falkowski, H.P. Nilles and M. Olechowski, “Soft supersymmetry breaking in KKLT flux compactification,” Nucl. Phys. B **718**, 113 (2005) [arXiv:hep-th/0503216].

[21] M.J. Duff, B.E.W. Nilsson and C.N. Pope, “Kaluza-Klein Supergravity,” Phys. Rept. **130**, 1 (1986).

[22] B. de Carlos, S. Gurrieri, A. Lukas and A. Micu, “Moduli stabilisation in heterotic string compactifications,” JHEP **0603**, 005 (2006) [arXiv:hep-th/0507173].

[23] D. Lüst and S. Stieberger, “Gauge threshold corrections in intersecting brane world models,” arXiv:hep-th/0302221.

[24] K. Choi, A. Falkowski, H.P. Nilles, M. Olechowski and S. Pokorski, “Stability of flux compactifications and the pattern of supersymmetry breaking,” JHEP **0411**, 076 (2004) [arXiv:hep-th/0411066].

[25] S. P. de Alwis, “Effective potentials for light moduli,” Phys. Lett. B **626**, 223 (2005) [arXiv:hep-th/0506266].

[26] H. Abe, T. Higaki and T. Kobayashi, “Remark on integrating out heavy moduli in flux compactification,” Phys. Rev. D **74**, 045012 (2006) [arXiv:hep-th/0606095];
H.X. Yang, “On moduli stabilization scheme in type IIB flux compactifications,” arXiv:hep-th/0608155; “Moduli stabilization in type IIB flux compactifications,” Phys. Rev. D **73**, 066006 (2006) [arXiv:hep-th/0511030].

[27] R.A. Horn and C.R. Johnson “Matrix Analysis”, Cambridge University Press, 1990.

[28] R. D’Auria, S. Ferrara and M. Trigiante, “c-map,very special quaternionic geometry and dual Kähler spaces,” Phys. Lett. B **587**, 138 (2004) [arXiv:hep-th/0401161].

[29] P. Candelas and X. de la Ossa, “Moduli Space Of Calabi-Yau Manifolds,” Nucl. Phys. B **355**, 455 (1991).

[30] J.J. Blanco-Pillado, C.P. Burgess, J.M. Cline, C. Escoda, M. Gomez-Reino, R. Kallosh, A. Linde and F. Quevedo, “Inflating in a better racetrack,” arXiv:hep-th/0603129.

[31] J.P. Conlon, “The QCD axion and moduli stabilisation,” arXiv:hep-th/0602233.

[32] M. Marino, R. Minasian, G.W. Moore and A. Strominger, "Nonlinear instantons from supersymmetric p-branes," JHEP **0001**, 005 (2000) [arXiv:hep-th/9911206].

[33] H. Jockers and J. Louis, "The effective action of D7-branes in N = 1 Calabi-Yau orientifolds," Nucl. Phys. B **705**, 167 (2005) [arXiv:hep-th/0409098]; "D-terms and F-terms from D7-brane fluxes," Nucl. Phys. B **718**, 203 (2005) [arXiv:hep-th/0502059]; H. Jockers, "The effective action of D-branes in Calabi-Yau orientifold compactifications," Fortsch. Phys. **53**, 1087 (2005) [arXiv:hep-th/0507042].

[34] M. Dine, N. Seiberg and E. Witten, "Fayet-Iliopoulos Terms In String Theory," Nucl. Phys. B **289**, 589 (1987).

[35] L.E. Ibanez, F. Marchesano and R. Rabadan, "Getting just the standard model at intersecting branes," JHEP **0111**, 002 (2001) [arXiv:hep-th/0105155].

[36] I. Antoniadis and T. Maillard, "Moduli stabilization from magnetic fluxes in type I string theory," Nucl. Phys. B **716**, 3 (2005) [arXiv:hep-th/0412008].

[37] A.K. Kashani-Poor and A. Tomasiello, "A stringy test of flux-induced isometry gauging," Nucl. Phys. B **728**, 135 (2005) [arXiv:hep-th/0505208].

[38] R. Kallosh and D. Sorokin, "Dirac action on M5 and M2 branes with bulk fluxes," JHEP **0505**, 005 (2005) [arXiv:hep-th/0501081].

[39] N. Saulina, "Topological constraints on stabilized flux vacua," Nucl. Phys. B **720**, 203 (2005) [arXiv:hep-th/0503125].

[40] R. Kallosh, A.K. Kashani-Poor and A. Tomasiello, "Counting fermionic zero modes on M5 with fluxes," JHEP **0506**, 069 (2005) [arXiv:hep-th/0503138].

[41] E. Bergshoeff, R. Kallosh, A.K. Kashani-Poor, D. Sorokin and A. Tomasiello, "An index for the Dirac operator on D3 branes with background fluxes," JHEP **0510**, 102 (2005) [arXiv:hep-th/0507069].

[42] J. Park, "D3 instantons in Calabi-Yau orientifolds with(out) fluxes," arXiv:hep-th/0507091.

[43] D. Lüst, S. Reffert, W. Schulgin and P.K. Tripathy, "Fermion zero modes in the presence of fluxes and a non-perturbative superpotential," arXiv:hep-th/0509082.

[44] L. Martucci, J. Rosseel, D. Van den Bleeken and A. Van Proeyen, "Dirac actions for D-branes on backgrounds with fluxes," Class. Quant. Grav. **22**, 2745 (2005) [arXiv:hep-th/0504041].

[45] L. Anguelova and K. Zoubos, "Five-brane instantons vs flux-induced gauging of isometries," arXiv:hep-th/0606271.

[46] I. Bandos and D. Sorokin, "Aspects of D-brane dynamics in supergravity backgrounds with fluxes, kappa-symmetry and equations of motion. IIB," arXiv:hep-th/0607163.

[47] Work in progress.

[48] L.J. Dixon, J.A. Harvey, C. Vafa and E. Witten, "Strings On Orbifolds," Nucl. Phys. B **261**, 678 (1985); "Strings On Orbifolds. 2," Nucl. Phys. B **274**, 285 (1986).

[49] D. Lüst, S. Reffert and S. Stieberger, “Flux-induced soft supersymmetry breaking in chiral type IIb orientifolds with D3/D7-branes,” Nucl. Phys. B **706**, 3 (2005) [arXiv:hep-th/0406092]

[50] G. Aldazabal, A. Font, L.E. Ibanez and G. Violero, “D = 4, N = 1, type IIB orientifolds,” Nucl. Phys. B **536**, 29 (1998) [arXiv:hep-th/9804026].

[51] G. Zwart, “Four-dimensional N = 1 Z(N) x Z(M) orientifolds,” Nucl. Phys. B **526**, 378 (1998) [arXiv:hep-th/9708040].

[52] M. Klein and R. Rabadan, “Z(N) x Z(M) orientifolds with and without discrete torsion,” JHEP **0010**, 049 (2000) [arXiv:hep-th/0008173]; “D = 4, N = 1 orientifolds with vector structure,” Nucl. Phys. B **596**, 197 (2001) [arXiv:hep-th/0007087]; “Orientifolds with discrete torsion,” JHEP **0007**, 040 (2000) [arXiv:hep-th/0002103].

[53] O. DeWolfe, A. Giryavets, S. Kachru and W. Taylor, “Enumerating flux vacua with enhanced symmetries,” JHEP **0502**, 037 (2005) [arXiv:hep-th/0411061].

[54] P. Berglund and P. Mayr, “Non-perturbative superpotentials in F-theory and string duality,” arXiv:hep-th/0504058.

[55] P.S. Aspinwall and R. Kallosh, “Fixing all moduli for M-theory on K3 x K3,” JHEP **0510**, 001 (2005) [arXiv:hep-th/0506014].

[56] P.G. Camara, L.E. Ibanez and A.M. Uranga, “Flux-induced SUSY-breaking soft terms on D7-D3 brane systems,” Nucl. Phys. B **708**, 268 (2005) [arXiv:hep-th/0408036].

[57] J. Gomis, F. Marchesano and D. Mateos, “An open string landscape,” JHEP **0511**, 021 (2005) [arXiv:hep-th/0506179].

[58] P. Mayr and S. Stieberger, “Threshold corrections to gauge couplings in orbifold compactifications,” Nucl. Phys. B **407**, 725 (1993) [arXiv:hep-th/9303017].